arXiv:1003.2017v2 [math.QA] 2 Jul 2010

THE TRIGONOMETRIC CASIMIR CONNECTION OF A SIMPLE LIE ALGEBRA

VALERIO TOLEDANO LAREDO

To Corrado De Concini, on his 60th birthday

Abstract. Let $\mathfrak{g}$ be a complex, semisimple Lie algebra, G the corresponding simply–connected Lie group and $H \subset G$ a maximal torus. We construct a flat connection on H with logarithmic singularities on the root hypertori and values in the Yangian $Y(\mathfrak{g})$ of $\mathfrak{g}$. By analogy with the rational Casimir connection of $\mathfrak{g}$, we conjecture that the monodromy of this trigonometric connection is described by the quantum Weyl group operators of the quantum loop algebra $U_\hbar(L\mathfrak{g})$.

Contents

1. Introduction 1
2. The trigonometric connection of a root system 4
3. The trigonometric Casimir connection 17
4. The monodromy conjecture 21
5. The trigonometric Casimir connection of $\mathfrak{gl}_n$ 22
6. Bispectrality 30
7. The affine KZ connection 34
8. Appendix: Tits extensions of affine Weyl groups 36

Acknowledgments 43

References 43

1. Introduction

1.1. Let $\mathfrak{g}$ be a complex, simple Lie algebra, $\mathfrak{h} \subset \mathfrak{g}$ a Cartan subalgebra, $\Phi \subset \mathfrak{h}^*$ the corresponding root system and W its Weyl group. For each $\alpha \in \Phi$, let $\mathfrak{sl}_2^\alpha = \langle e_\alpha, f_\alpha, h_\alpha \rangle \subset \mathfrak{g}$ be the corresponding three–dimensional subalgebra and denote by

$$\kappa_\alpha = \frac{(\alpha,\alpha)}{2}\left(e_\alpha f_\alpha + f_\alpha e_\alpha\right)$$

Supported in part by NSF grants DMS–0707212 and DMS–0854792.

its truncated Casimir operator with respect to the restriction to $\mathfrak{sl}_2^\alpha$ of a fixed non–degenerate, ad–invariant bilinear form $(\cdot,\cdot)$ on $\mathfrak{g}$. Let

$$\mathfrak{h}_{\text{reg}} = \mathfrak{h} \setminus \bigcup_{\alpha\in\Phi} \operatorname{Ker}(\alpha)$$

be the set of regular elements in $\mathfrak{h}$, V a finite–dimensional $\mathfrak{g}$–module, and $\mathbb{V}$ the holomorphically trivial vector bundle over $\mathfrak{h}_{\text{reg}}$ with fibre V. Recall that the Casimir connection of $\mathfrak{g}$ is the holomorphic connection on $\mathbb{V}$ given by

$$\nabla_\kappa = d - \hbar \sum_{\alpha\in\Phi_+} \frac{d\alpha}{\alpha}\,\kappa_\alpha \tag{1.1}$$

where d is the de Rham differential, $\Phi_+ \subset \Phi$ is a system of positive roots and $\hbar$ is a complex number. This connection was discovered independently by C. De Concini around 1995 (unpublished), by J. Millson and the author [MTL, TL1] and Felder *et al.* [FMTV], and shown to be flat for any $\hbar \in \mathbb{C}$.

The monodromy of ∇_κ gives representations of the generalised braid group $B = \pi_1(\mathfrak{h}_{\text{reg}}/W)$ which are described by the quantum Weyl group operators of the quantum group $U_\hbar\mathfrak{g}$, a fact which was conjectured by De Concini (unpublished) and independently in [TL1, TL2] and proved in [TL1, TL3].

1.2. Let $P \subset \mathfrak{h}^*$ be the weight lattice of $\mathfrak{g}$ and $H = \operatorname{Hom}_{\mathbb{Z}}(P, \mathbb{C}^*)$ the dual algebraic torus with Lie algebra $\mathfrak{h}$ and coordinate ring given by the group algebra $\mathbb{C}P$. We denote the function corresponding to $\lambda \in P$ by $e^\lambda \in \mathbb{C}[H]$. The main goal of the present paper is to define a *trigonometric* version of the connection (1.1), that is a connection defined on H with the logarithmic forms $d\alpha/\alpha$ replaced by $d\alpha/(e^\alpha - 1)$.

As is well known from the study of Cherednik's affine KZ (AKZ) connection (see, *e.g.* [Ch3]), both the flatness and W–equivariance of such a connection require that it possess a 'tail', that is be of the form

$$\widehat{\nabla}_\kappa = d - \hbar \sum_{\alpha\in\Phi_+} \frac{d\alpha}{e^\alpha - 1}\,\kappa_\alpha - A \tag{1.2}$$

where A is a translation–invariant one–form on H. The analogy with the AKZ equations further suggests that A should take values in a suitable extension of the enveloping algebra $U\mathfrak{g}$, which is to $U\mathfrak{g}$ what the degenerate affine Hecke algebra $\mathcal{H}'$ is to the group algebra $\mathbb{C}W$.

1.3. The correct extension turns out to be the *Yangian* $Y(\mathfrak{g})$, which is a deformation of the enveloping algebra $U(\mathfrak{g}[t])$ over the ring $\mathbb{C}[\hbar]$. Let $\nu : \mathfrak{h} \to \mathfrak{h}^*$ be the isomorphism determined by the inner product $(\cdot,\cdot)$, set $t_i = \nu^{-1}(\alpha_i)$, where $\alpha_1, \ldots, \alpha_n$ are the simple roots of $\mathfrak{g}$ relative to Φ_+ and let $t^i = \lambda_i^\vee$ be the dual basis of $\mathfrak{h}$ given by the fundamental coweights. Let $T(u)_r$, $u \in \mathfrak{h}$, $r \in \mathbb{N}$ be the Cartan loop generators of $Y(\mathfrak{g})$ in Drinfeld's new realisation (see [Dr2] and §3 for definitions). Let $\{u^i\}$ be a basis of $\mathfrak{h}$, $\{u_i\}$ the dual basis of $\mathfrak{h}^*$ and regard the differentials du_i as translation–invariant one–forms on H. The main result of this paper is the following

Theorem. *The $Y(\mathfrak{g})$–valued connection on H given by*

$$\widehat{\nabla}_\kappa = d - \hbar \sum_{\alpha\in\Phi_+} \frac{d\alpha}{e^\alpha - 1}\,\kappa_\alpha + du_i\left(2T(u^i)_1 - \frac{\hbar}{2}(u^i,t^j)t_j^2\right)$$

is flat and W–equivariant[1].

1.4. We call $\widehat{\nabla}_\kappa$ the *trigonometric Casimir connection* of $\mathfrak{g}$. Its monodromy defines representations of the *affine braid group* $\widehat{B} = \pi_1(H_{\text{reg}}/W)$ on any finite–dimensional module over $Y(\mathfrak{g})$, where

$$H_{\text{reg}} = H \setminus \bigcup_{\alpha\in\Phi}\{e^\alpha = 1\}$$

is the set of regular elements in H. By analogy with the rational case, we conjecture that these representations are equivalent to the quantum Weyl group action of $\widehat{B}$ on finite–dimensional modules over the quantum loop algebra $U_\hbar(\mathfrak{g}[z,z^{-1}])$.

1.5. We turn now to a detailed description of the paper.

In Section 2 we obtain a necessary and sufficient condition for a connection of the form (1.2) to be flat and equivariant under the Weyl group W (Theorem 2.5 and Proposition 2.22), thus effectively determining the Lie algebras of the fundamental groups $\pi_1(H_{\text{reg}})$ and $\pi_1(H_{\text{reg}}/W)$. Consistently with Cherednik's study of the AKZ connection, the quadratic equations giving the flatness and equivariance of $\widehat{\nabla}_\kappa$ specialise, when κ_α is replaced by the orthogonal reflection $s_\alpha \in W$ to the defining relations of the degenerate affine Hecke algebra corresponding to W.

In Section 3 we review Drinfeld's two presentations of the Yangian $Y(\mathfrak{g})$ of a simple Lie algebra $\mathfrak{g}$. We then use their interplay to solve the above quadratic equations in $Y(\mathfrak{g})$, thereby obtaining the trigonometric Casimir connection of $\mathfrak{g}$ (Theorem 3.8).

In Section 4, we explain how to define monodromy representations of the affine braid group from $\widehat{\nabla}_\kappa$ and conjecture that these are described by the quantum Weyl group operators of the quantum loop algebra $U_\hbar(\mathfrak{g}[z,z^{-1}])$.

In Section 5, we define a trigonometric connection with values in the Yangian of $\mathfrak{gl}_n$ by using the interplay between its loop and RTT presentations (Theorem 5.7). We then relate it to the trigonometric Casimir connection of $\mathfrak{sl}_n$. We also check that, when computed in a tensor product of m evaluation modules, it coincides with the trigonometric dynamical differential equations [TV] which are differential equations on $(\mathbb{C}^\times)^n$ with values in $U\mathfrak{gl}_n^{\otimes m}$.

In Section 6, we show that the trigonometric Casimir connection commutes with qKZ difference equations of Frenkel–Reshetikhin determined by the rational R–matrix of $Y(\mathfrak{g})$ (Theorem 6.6), a fact which was checked in [TV] for the trigonometric dynamical differential equations.

---

[1] we follow the standard convention that in any expression involving u_i and u^i, or t^i and t_i, summation over i is implicit.

In Section 7, we review the definition of the degenerate affine Hecke algebra $\mathcal{H}'$ of W [Lu] and of the corresponding $\mathcal{H}'$–valued AKZ connection [Ch3]. We then show that if V is a $Y(\mathfrak{g})$–module whose restriction to $\mathfrak{g}$ is *small*, that is such that 2α is not a weight for any root α [Br, Re], the zero weight space $V[0]$ carries a natural action of $\mathcal{H}'$. Moreover, the trigonometric Casimir connection with coefficients in $V[0]$ coincides with the AKZ connection with values in this $\mathcal{H}'$–module (Theorem 7.5).

The final appendix, Section 8 contains a discussion of the Tits extensions of affine Weyl groups which is needed for Section 4.

2. The trigonometric connection of a root system

2.1. **General form.** Let E be a Euclidean vector space, $\Phi \subset E^*$ a reduced, crystallographic root system. Let $Q^\vee \subset E$ be the lattice generated by the coroots $\alpha^\vee$, $\alpha \in \Phi$ and $P \subset E^*$ the dual weight lattice. Let $H = \mathrm{Hom}_{\mathbb{Z}}(P, \mathbb{C}^*)$ be the complex algebraic torus with Lie algebra $\mathfrak{h} = \mathrm{Hom}_{\mathbb{Z}}(P, \mathbb{C})$ and coordinate ring given by the group algebra $\mathbb{C}P$. We denote the function corresponding to $\lambda \in P$ by $e^\lambda \in \mathbb{C}[H]$ and set

$$H_{\mathrm{reg}} = H \setminus \bigcup_{\alpha \in \Phi} \{e^\alpha = 1\} \tag{2.1}$$

Let A be an algebra endowed with the following data:

- a set of elements $\{t_\alpha\}_{\alpha \in \Phi} \subset A$ such that $t_{-\alpha} = t_\alpha$
- a linear map $\tau : \mathfrak{h} \to A$

Consider the A–valued connection on H_{reg} given by

$$\nabla = d - \sum_{\alpha \in \Phi_+} \frac{d\alpha}{e^\alpha - 1}\, t_\alpha - du_i\, \tau(u^i) \tag{2.2}$$

where $\Phi_+ \subset \Phi$ is a chosen system of positive roots, $\{u_i\}$ and $\{u^i\}$ are dual bases of $\mathfrak{h}^*$ and $\mathfrak{h}$ respectively, the differentials du_i are regarded as translation–invariant one–forms on H and the summation over i is implicit.

2.2. **Positive roots.** The form of the connection (2.2) depends upon the choice of the system of positive roots $\Phi_+ \subset \Phi$. Let however $\Phi'_+ \subset \Phi$ be another such system, then

Proposition. *The connection* (2.2) *may be rewritten as*

$$\nabla = d - \sum_{\alpha \in \Phi'_+} \frac{d\alpha}{e^\alpha - 1}\, t_\alpha - du_i\, \tau'(u^i)$$

where $\tau' : \mathfrak{h} \to A$ *is given by*

$$\tau'(v) = \tau(v) - \sum_{\alpha \in \Phi_+ \cap \Phi'_-} \alpha(v)\, t_\alpha \tag{2.3}$$

Proof. Write the second summand in (2.2) as

$$\sum_{\alpha\in\Phi_+\cap\Phi'_+}\frac{d\alpha}{e^{\alpha}-1}t_{\alpha}-\sum_{\alpha\in\Phi_-\cap\Phi'_+}\frac{d\alpha}{e^{-\alpha}-1}t_{-\alpha}$$

where $\Phi_-=-\Phi_+$. Since

$$\frac{1}{1-e^{-\alpha}}=\frac{e^{\alpha}}{e^{\alpha}-1}=\frac{1}{e^{\alpha}-1}+1 \tag{2.4}$$

and $t_{-\alpha}=t_{\alpha}$, the above is equal to

$$\sum_{\alpha\in\Phi'_+}\frac{d\alpha}{e^{\alpha}-1}t_{\alpha}+\sum_{\alpha\in\Phi_-\cap\Phi'_+}d\alpha\,t_{\alpha}$$

which yields the required result since $\alpha=u_i\alpha(u^i)$. ■

Definition. If W is the Weyl group of Φ and $w\in W$ the unique element such that $\Phi'_+=w\Phi_+$, we denote τ' by τ_w. Thus,

$$\tau_w(v)=\tau(v)-\sum_{\alpha\in\Phi_+\cap w\Phi_-}\alpha(v)\,t_{\alpha} \tag{2.5}$$

2.3. **Delta form.** Choose $\Phi'_+=\Phi_-$ in Proposition 2.2. Comparing the corresponding expressions for ∇ shows that it may be more invariantly rewritten as

$$\nabla=d-\frac{1}{2}\sum_{\alpha\in\Phi}\frac{d\alpha}{e^{\alpha}-1}t_{\alpha}-du_i\,\delta(u^i)$$

where $\delta:\mathfrak{h}\to A$ is given by

$$\delta(v)=\tau(v)-\frac{1}{2}\sum_{\alpha\in\Phi_+}\alpha(v)\,t_{\alpha} \tag{2.6}$$

Alternatively, substituting (2.6) into (2.2) yields

$$\nabla=d-\frac{1}{2}\sum_{\alpha\in\Phi_+}\frac{e^{\alpha}+1}{e^{\alpha}-1}d\alpha\,t_{\alpha}-du_i\,\delta(u^i) \tag{2.7}$$

We shall occasionally refer to (2.2) and (2.7) as the τ *and* δ*–forms* of the connection ∇ respectively. Note that the latter does not depend upon the choice of $\Phi_+\subset\Phi$.

2.4. **Root subsystems.** For a subset $\Psi\subset\Phi$ and subring $R\subset\mathbb{R}$, let $\langle\Psi\rangle_R\subset E^*$ be the R–span of Ψ.

Definition. A *root subsystem* of Φ is a subset $\Psi\subset\Phi$ such that $\langle\Psi\rangle_{\mathbb{Z}}\cap\Phi=\Psi$. Ψ is *complete* if $\langle\Psi\rangle_{\mathbb{R}}\cap\Phi=\Psi$. If $\Psi\subset\Phi$ is a root subsystem, we set $\Psi_+=\Psi\cap\Phi_+$.

Remark. According to the above definition, the short roots of the root system B_2 (resp. G_2) are not a root subsystem, but the long ones constitute a root subsystem of type $\mathsf{A}_1 \times \mathsf{A}_1$ (resp. A_2) which is not complete. Another root subsystem of $\Phi = \mathsf{G}_2$ is given by $\{\pm\alpha, \pm\beta\}$ where α, β are two orthogonal roots (necessarily of different lengths).

2.5. **Integrability.** The following is the main result of this section.

Theorem.

(1) *The connection* ∇ *is flat if, and only if the following relations hold*

- *For any rank 2 root subsystem* $\Psi \subset \Phi$ *and* $\alpha \in \Psi$,
$$[t_\alpha, \sum_{\beta \in \Psi_+} t_\beta] = 0 \tag{tt}$$
- *For any* $u, v \in \mathfrak{h}$,
$$[\tau(u), \tau(v)] = 0 \tag{$\tau\tau$}$$
- *For any* $\alpha \in \Phi_+$, $w \in W$ *such that* $w^{-1}\alpha$ *is a simple root and* $u \in \mathfrak{h}$ *such that* $\alpha(u) = 0$,
$$[t_\alpha, \tau_w(u)] = 0 \tag{$t\tau$}$$

(2) *Modulo the relations* (tt), *the relations* $(t\tau)$ *are equivalent to*
$$[t_\alpha, \delta(v)] = 0 \tag{$t\delta$}$$
for any $\alpha \in \Phi$ *and* $v \in \mathfrak{h}$ *such that* $\alpha(v) = 0$, *where* $\delta : \mathfrak{h} \to A$ *is given by* (2.6).

The proof of Theorem 2.5 occupies the paragraphs 2.7–2.19.

2.6. We spell out below the relations (tt) in the case when Φ is of rank 2. For $\Psi = \Phi$, they read
$$[t_\alpha, \sum_{\beta \in \Phi_+} t_\beta] = 0 \quad \text{for any } \alpha \in \Phi \tag{Φ}$$
In particular, if $\Phi = \{\pm\alpha, \pm\beta\}$ is of type $\mathsf{A}_1 \times \mathsf{A}_1$ then
$$[t_\alpha, t_\beta] = 0 \tag{$\mathsf{A}_1 \times \mathsf{A}_1$}$$
For $\Phi = \mathsf{B}_2$, the long roots $\{\pm\beta_1, \pm\beta_2\}$ form an $\mathsf{A}_1 \times \mathsf{A}_1$ subsystem so that
$$[t_{\beta_1}, t_{\beta_2}] = 0 \tag{$\mathsf{A}_1 \times \mathsf{A}_1 \subset \mathsf{B}_2$}$$
For $\Phi = \mathsf{G}_2$, there are two types of root subsystems: that formed by the A_2 configuration of long roots $\{\pm\beta_1, \pm\beta_2, \pm\beta_3\}$, leading to
$$[t_{\beta_i}, t_{\beta_j} + t_{\beta_k}] = 0 \quad \text{for any } i, j, k \in \{1, 2, 3\} \text{ distinct} \tag{$\mathsf{A}_2 \subset G_2$}$$
and the A_1 configurations $\{\pm\beta, \pm\gamma\}$ formed by a long root and an orthogonal short one, leading to
$$[t_\beta, t_\gamma] = 0 \tag{$\mathsf{A}_1 \times \mathsf{A}_1 \subset \mathsf{G}_2$}$$

Combining relations (Φ), $(\mathsf{A}_2 \subset G_2)$ and $(\mathsf{A}_1 \times \mathsf{A}_1 \subset \mathsf{G}_2)$ yields in particular the following relations

$$[t_\beta, t_{\gamma'} + t_{\gamma''}] = 0 \tag{2.8}$$

where $\beta \in \mathsf{G}_2$ is long and γ', γ'' are the short positive roots which are not orthogonal to β.

Remark. If Φ is not simply–laced, the relations (tt) are stronger than those yielding the flatness of the rational connection

$$\nabla = d - \sum_{\alpha \in \Phi_+} \frac{d\alpha}{\alpha} t_\alpha$$

Indeed, the latter involve two dimensional subspaces of $\mathfrak{h}^*$ spanned by elements of Φ [Ko] and therefore only those rank 2 subsystems of Φ which are complete. The relevance of additional relations corresponding to non–complete subsystems was first pointed out in the closely related context of the Yang–Baxter equations by Cherednik [Ch3, §6.1].

2.7. Let $Q \subset P$ be the root lattice generated by Φ and $T = \mathrm{Hom}_{\mathbb{Z}}(Q, \mathbb{C}^*)$ the corresponding complex algebraic torus of adjoint type. T has coordinate ring $\mathbb{C}Q$ and is biregularly isomorphic to the standard torus $(\mathbb{C}^*)^n$ by sending $p \in T$ to the point with coordinates $z_i = e^{-\alpha_i}(p)$, where α_i varies over the simple roots of Φ relative to Φ_+.

2.8. The torus T is a quotient of H and the form of ∇ shows that it may be regarded as a connection on the trivial vector bundle with fibre A over T. As such, ∇ has singularities on the codimension one subtori

$$T_\alpha = \{e^\alpha = 1\} \subset T$$

where $\alpha \in \Phi$. Given a subset $\Psi \subset \Phi$, we shall be interested in the connectedness of the intersection $\bigcap_{\alpha \in \Psi} T_\alpha$. Let $\langle \Psi \rangle_{\mathbb{Z}} \subset Q$ be the $\mathbb{Z}$–span of Ψ and set [DCP, §3.1]

$$\overline{\langle \Psi \rangle}_{\mathbb{Z}} = \{\gamma \in Q |\, m\gamma \in \langle \Psi \rangle_{\mathbb{Z}} \text{ for some } m \in \mathbb{Z}^*\}$$

Since

$$\mathbb{C}[\bigcap_{\alpha \in \Psi} T_\alpha] = \mathbb{C}\, Q / \langle \Psi \rangle_{\mathbb{Z}} \cong \mathbb{C}\, Q / \overline{\langle \Psi \rangle}_{\mathbb{Z}} \otimes \mathbb{C}\, \overline{\langle \Psi \rangle}_{\mathbb{Z}} / \langle \Psi \rangle_{\mathbb{Z}}$$

the connected components of $\bigcap_{\alpha \in \Psi} T_\alpha$ are tori labelled by the characters of the finite abelian group $\overline{\langle \Psi \rangle}_{\mathbb{Z}} / \langle \Psi \rangle_{\mathbb{Z}}$. In particular, if $\Psi = \{\alpha\}$, we see that each T_α is connected since α is indivisible in Q.

2.9. The necessity of relations (tt)–$(t\tau)$ follows from the computation of the residues of the curvature Ω of ∇ to be carried out in §2.10–§2.14.

Specifically, write $\nabla = d - A$. Since $dA = 0$, Ω is equal to $A\wedge A = \Omega_1+\Omega_2+\Omega_3$, where

$$\Omega_1 = \frac{1}{2}\sum_{\alpha,\beta}\frac{d\alpha}{e^\alpha-1}\wedge\frac{d\beta}{e^\beta-1}\,[t_\alpha,t_\beta] \tag{2.9}$$

$$\Omega_2 = \sum_{\alpha,i}\frac{d\alpha}{e^\alpha-1}\wedge du_i\,[t_\alpha,\tau(u^i)] \tag{2.10}$$

$$\Omega_3 = \frac{1}{2}\sum_{i,j} du_i\wedge du_j\,[\tau(u^i),\tau(u^j)] \tag{2.11}$$

2.10. Let $\alpha\in\Phi$ and denote the inclusion $T_\alpha\hookrightarrow T$ by $\imath_\alpha$. Then

$$\operatorname{res}_{T_\alpha}\Omega_1 = \imath_\alpha^*\sum_{\beta\neq\alpha}\frac{d\beta}{e^\beta-1}\,[t_\alpha,t_\beta]$$

$$\operatorname{res}_{T_\alpha}\Omega_2 = \imath_\alpha^* du_i\,[t_\alpha,\tau(u^i)]$$

and $\operatorname{res}_{T_\alpha}\Omega_3 = 0$ since Ω_3 is regular on T_α.

2.11. Let $\Psi\subset\Phi$ be a rank 2 root subsystem and set

$$T_\Psi = \bigcap_{\beta\in\Psi} T_\beta$$

By §2.8, T_Ψ is a codimension two subtorus of T with group of components $\operatorname{Hom}(\overline{\langle\Psi\rangle}_\mathbb{Z}/\langle\Psi\rangle_\mathbb{Z},\mathbb{C}^*)$.

Since $\overline{\langle\Psi\rangle}_\mathbb{Z}/\langle\Psi\rangle_\mathbb{Z}$ is cyclic (of order 1, 2 or 3 depending on the type of Ψ and $\overline{\langle\Psi\rangle}_\mathbb{Z}\cap\Phi$), there exists a character χ of $\overline{\langle\Psi\rangle}_\mathbb{Z}/\langle\Psi\rangle_\mathbb{Z}$ with trivial kernel. It follows that the corresponding component T_Ψ^χ of T_Ψ is contained in some T_γ if, and only if, $\gamma\in\Psi$.

Together with §2.10, this implies that for any $\alpha\in\Psi$,

$$\operatorname{res}_{T_\Psi^\gamma}\operatorname{res}_{T_\alpha}\Omega = [t_\alpha,\sum_{\beta\in\Psi_+}t_\beta]$$

thus showing the necessity of (tt).

2.12. Let $\overline{T}\cong\mathbb{C}^n$ be the partial compactification determined by the embedding $T\hookrightarrow(\mathbb{C}^*)^n$ given by sending $p\in T$ to the point with coordinates $z_i = e^{-\alpha_i}(p)$. We wish to determine the residues of Ω on the divisors

$$T_i = \{z_i = 0\}\subset\overline{T}$$

To this end, we first rewrite ∇ in the coordinates z_i. Choosing $u_i=\alpha_i$ as basis of $\mathfrak{h}^*$, so that the dual basis $\{u^i\}$ of $\mathfrak{h}$ is given by the fundamental coweights $\{\lambda_i^\vee\}$ yields $du_i = -dz_i/z_i$ and

$$du_i\,\tau(u^i) = -\frac{dz_i}{z_i}\,\tau(\lambda_i^\vee)$$

Further, if $\alpha = \sum_i m_\alpha^i \alpha_i$ is a positive root, then $e^\alpha = \prod_i z_i^{-m_\alpha^i}$ so that

$$\frac{d\alpha}{e^\alpha - 1} = \frac{e^{-\alpha}}{1 - e^{-\alpha}}\, d\alpha = -\sum_i m_\alpha^i \frac{z_i^{m_\alpha^i - 1} \prod_{j \neq i} z_j^{m_\alpha^j}}{1 - \prod_j z_j^{m_\alpha^j}}\, dz_i$$

which is a regular on each T_i. It follows that $\operatorname{res}_{T_i} \Omega_1 = 0$ and

$$\operatorname{res}_{T_i} \Omega_2 = \imath_i^* \sum_\alpha \frac{d\alpha}{e^\alpha - 1} [t_\alpha, \tau(\lambda_i^\vee)]$$

$$\operatorname{res}_{T_i} \Omega_3 = \imath_i^* \sum_{j \neq i} \frac{dz_j}{z_j} [\tau(\lambda_i^\vee), \tau(\lambda_j^\vee)]$$

where $\imath_i$ is the inclusion $T_i \hookrightarrow \overline{T}$.

2.13. Thus, for any $j \neq i$,

$$\operatorname{res}_{T_j \cap T_i} \operatorname{res}_{T_i} \Omega = [\tau(\lambda_j^\vee), \tau(\lambda_i^\vee)]$$

which shows the necessity of the relations $(\tau\tau)$.

2.14. Let now $\alpha = \sum_i m_\alpha^i \alpha_i$ be a positive root and let $\overline{T}_\alpha = \{\prod_i z_i^{m_\alpha^i} = 1\}$ be the closure of T_α in $\overline{T}$. The intersection

$$T_{\alpha,i} = \overline{T}_\alpha \cap T_i \subset \overline{T}$$

is clearly nonempty if, and only if $\alpha(\lambda_i^\vee) = m_\alpha^i = 0$. When that is the case, $T_{\alpha,i}$ is connected and contained in no other $\overline{T}_\beta$, $\beta \neq j$ or T_j for $j \neq i$.

It follows that whenever $\alpha(\lambda_i^\vee) = 0$,

$$\operatorname{res}_{T_{\alpha,i}} \operatorname{res}_{T_\alpha} \Omega = [t_\alpha, \tau(\lambda_i^\vee)]$$

thus showing the necessity of $(t\tau)$ for α simple and $w = 1$. The general case follows by repeating the computations of the last two subsections in the compactification of T corresponding to a different basis $w\Delta$ of simple roots and using the alternative form of the connection ∇ given by Proposition 2.2.

2.15. We next turn to the sufficiency of the relations (tt)–$(\tau\tau)$. This may be proved by embedding T in the toric variety corresponding to the fan determined by the chambers of Φ in E and using a general integrability criterion of E. Looijenga as in [Lo, §1–2][2]. We prefer a more direct approach which will occupy §2.16– 2.19.

---

[2] note however that line 2 of the statement of Corollary 1.3 in [Lo] should read "for every **irreducible component** I **of a** codimension two intersection", the words in bold are missing in [Lo].

2.16. Since the relations $(\tau\tau)$ imply that $\Omega_3 = 0$, we need to show that the relations (tt)–$(t\tau)$ imply that $\Omega_1 + \Omega_2 = 0$. To this end, we rewrite Ω_2 in a different form below and, in §2.17 rewrite Ω_1.

Lemma. *Modulo the relations $(t\tau)$, the curvature term*

$$\Omega_2^\alpha = \sum_i \frac{d\alpha}{e^\alpha - 1} \wedge du_i\,[t_\alpha, \tau(u^i)]$$

corresponding to $\alpha \in \Phi_+$ is equal to

$$\sum_{\beta \in \Phi_+ \cap w\Phi_-} \frac{d\alpha}{e^\alpha - 1} \wedge d\beta\,[t_\alpha, t_\beta]$$

for any $w \in W$ such that $w^{-1}\alpha$ is a simple root.

Proof. Let $w \in W$ be such that $w^{-1}\alpha$ is a simple root α_i. By (2.5),

$$\Omega_2^\alpha = \sum_j \frac{d\alpha}{e^\alpha - 1} \wedge du_j\,[t_\alpha, \tau_w(u^j) + \sum_{\beta \in \Phi_+ \cap w\Phi_-} \beta(u^j)\,t_\beta]$$

Choosing $u_j = w\alpha_j$ yields a commutator $[t_\alpha, \tau_w(u^j)] = [t_\alpha, \tau_w(w\lambda_j^\vee)]$ which is zero for all $j \neq i$ by $(t\tau)$. Since $d\alpha \wedge w\alpha_i = 0$, this yields

$$\Omega_2^\alpha = \sum_j \frac{d\alpha}{e^\alpha - 1} \wedge du_j\,[t_\alpha, \sum_{\beta \in \Phi_+ \cap w\Phi_-} \beta(u^j)\,t_\beta] = \sum_{\beta \in \Phi_+ \cap w\Phi_-} \frac{d\alpha}{e^\alpha - 1} \wedge d\beta\,[t_\alpha, t_\beta]$$

since $\beta = \beta(u^i)u_i$ ■

2.17. For any $a \in Q$, let η_a be the meromorphic one–form on T given by

$$\eta_a = \frac{da}{e^a - 1}$$

Lemma. *Let $\Psi \subset \Phi$ be a rank 2 root subsystem and consider the curvature term*

$$\Omega_1^\Psi = \frac{1}{2} \sum_{\alpha, \beta \in \Psi_+} \eta_\alpha \wedge \eta_\beta\,[t_\alpha, t_\beta]$$

Assume that the relations (tt) hold. Then, if Ψ is of type $\mathsf{A}_1 \times \mathsf{A}_1$,

$$\Omega_1^\Psi = 0 \tag{2.12}$$

If Ψ is of type A_2 with $\Psi_+ = \{\alpha, \beta, \alpha + \beta\}$

$$\Omega_1^\Psi = -\eta_{\alpha+\beta} \wedge d\beta\,[t_{\alpha+\beta}, t_\beta] \tag{2.13}$$

If Ψ is of type B_2 with $\Psi_+ = \{\alpha, \beta, \alpha \pm \beta\}$

$$\Omega_1^\Psi = -\eta_{\alpha+\beta} \wedge d\beta\,[t_{\alpha+\beta}, t_\beta] - \eta_\alpha \wedge d(\alpha - \beta)\,[t_\alpha, t_{\alpha-\beta}] \tag{2.14}$$

If Ψ is of type G_2, with $\Psi_+ = \{\alpha_1, \alpha_2, \alpha_1+\alpha_2, \alpha_1+2\alpha_2, \alpha_1+3\alpha_2, 2\alpha_1+3\alpha_2\}$

$$\begin{aligned}\Omega_1^\Psi = &- \eta_{\alpha_1+\alpha_2} \wedge d\alpha_1\,[t_{\alpha_1+\alpha_2}, t_{\alpha_1}]\\ &- \eta_{\alpha_1+2\alpha_2} \wedge (d\alpha_2\,[t_{\alpha_1+2\alpha_2}, t_{\alpha_2}] + d(\alpha_1+3\alpha_2)\,[t_{\alpha_1+2\alpha_2}, t_{\alpha_1+3\alpha_2}])\\ &- \eta_{\alpha_1+3\alpha_2} \wedge d\alpha_2\,[t_{\alpha_1+3\alpha_2}, t_{\alpha_2}]\\ &- \eta_{2\alpha_1+3\alpha_2} \wedge (d\alpha_1\,[t_{2\alpha_1+3\alpha_2}, t_{\alpha_1}] + d(\alpha_1+\alpha_2)\,[t_{2\alpha_1+3\alpha_2}, t_{\alpha_1+\alpha_2}])\end{aligned} \tag{2.15}$$

Proof. If Ψ is of type $\mathsf{A}_1 \times \mathsf{A}_1$, the result follows from ($\mathsf{A}_1 \times \mathsf{A}_1$).

For other types, we shall need the following easily verified identity. For $a, b \in Q$, set

$$\eta_{a,b} = \frac{da \wedge db}{e^{a+b} - 1}$$

Then,

$$\eta_a \wedge \eta_b = \eta_a \wedge \eta_{a+b} + \eta_{a+b} \wedge \eta_b + \eta_{a,b} \tag{2.16}$$

We shall apply (2.16) to Ω_1^Ψ repeatedly, specifically to terms of the form $\eta_\alpha \wedge \eta_\beta$ with $\alpha + \beta \in \Phi_+$, until no such terms are left.

For $\Psi_+ = \{\alpha, \beta, \alpha+\beta\}$ of type A_2, this yields

$$\Omega_1^\Psi = \eta_{\alpha+\beta} \wedge \eta_\alpha\,[t_{\alpha+\beta} + t_\beta, t_\alpha] + \eta_{\alpha+\beta} \wedge \eta_\beta\,[t_{\alpha+\beta} + t_\alpha, t_\beta] + \eta_{\alpha,\beta}\,[t_\alpha, t_\beta]$$

By (Φ), the first two commutators are 0 and the third is equal to $[t_\alpha, t_\beta] = [t_\alpha + t_\beta, t_\beta] = -[t_{\alpha+\beta}, t_\beta]$. This yields the required answer since

$$\eta_{a,b} = \eta_{a+b} \wedge db = -\eta_{a+b} \wedge da \tag{2.17}$$

To keep track of the repeated applications of (2.16) for Ψ ot type $\mathsf{B}_2, \mathsf{G}_2$, we proceed as follows. Recall that the height of $\alpha = \sum_i m_\alpha^i \alpha_i \in \Phi_+$ is defined by $\mathrm{ht}(\alpha) = \sum_i m_\alpha^i$. Arrange pairs of distinct roots (α, β) on consecutive rows according to the value of $\mathrm{ht}(\alpha) + \mathrm{ht}(\beta)$: each (α, β) stands for a term $\eta_\alpha \wedge \eta_\beta$. From each pair (α, β) such that $\alpha + \beta \in \Phi_+$ draw an arrow to $(\alpha, \alpha+\beta)$ and $(\alpha+\beta, \beta)$ to signify that (2.16) has been applied with $a = \alpha$ and $b = \beta$.

For $\Psi_+ = \{\alpha, \beta, \alpha \pm \beta\}$ of type B_2 with simple roots $\alpha_1 = \alpha - \beta$, $\alpha_2 = \beta$, the corresponding graph is

$$\begin{tikzcd}
 & & (\alpha-\beta,\beta) \arrow[dl] \arrow[dr] & & \\
 & (\alpha-\beta,\alpha) & & (\alpha,\beta) \arrow[dr] \arrow[ddl] & \\
(\alpha-\beta,\alpha+\beta) & & & & (\alpha+\beta,\beta) \\
 & & (\alpha,\alpha+\beta) & &
\end{tikzcd}$$

This yields an "$\eta_a \wedge \eta_b$" component of Ω_1^Ψ equal to

$$\begin{aligned}\eta_{\alpha-\beta}\wedge\eta_\alpha\,[t_{\alpha-\beta},t_\alpha+t_\beta]&\\ +\eta_{\alpha-\beta}\wedge\eta_{\alpha+\beta}\,[t_{\alpha-\beta},t_{\alpha+\beta}]+\eta_{\alpha+\beta}\wedge\eta_\beta\,[t_{\alpha+\beta}+t_\alpha+t_{\alpha-\beta},t_\beta]&\\ +\eta_\alpha\wedge\eta_{\alpha+\beta}\,\left([t_\alpha,t_{\alpha+\beta}+t_\beta]+[t_{\alpha-\beta},t_\beta]\right)&\end{aligned}$$

The second commutator is equal to zero by ($\mathsf{A}_1\times\mathsf{A}_1\subset\mathsf{B}_2$), the first by ($\Phi$) and ($\mathsf{A}_1\times\mathsf{A}_1\subset\mathsf{B}_2$) and the third by ($\Phi$). By ($\Phi$), the coefficient of $\eta_\alpha\wedge\eta_{\alpha+\beta}$ is equal to $-[t_\alpha,t_{\alpha-\beta}]+[t_{\alpha-\beta},t_\beta]=[t_{\alpha-\beta},t_\alpha+t_\beta]=0$.

Ω_1^Ψ is therefore equal to its "$\eta_{a,b}$" component, namely

$$\eta_{\alpha-\beta,\beta}\,[t_{\alpha-\beta},t_\beta]+\eta_{\alpha,\beta}\,[t_\alpha+t_{\alpha-\beta},t_\beta]$$

which yields the required answer since, by (Φ)

$$[t_\alpha+t_{\alpha-\beta},t_\beta]=-[t_{\alpha+\beta},t_\beta]$$

and by (Φ) and ($\mathsf{A}_1\times\mathsf{A}_1\subset\mathsf{B}_2$)

$$[t_{\alpha-\beta},t_\beta]=-[t_{\alpha-\beta},t_\alpha+t_{\alpha+\beta}]=-[t_{\alpha-\beta},t_\alpha]$$

while, as previously noted

$$\eta_{a,b}=\eta_{a+b}\wedge db=-\eta_{a+b}\wedge da$$

Assume now that Ψ is of type G_2 and has simple roots α_1,α_2, with α_1 long. The sets of long and short positive roots are, respectively

$$\Psi_+^\ell=\{\alpha_1,2\alpha_1+3\alpha_2,\alpha_1+3\alpha_2\}\qquad\text{and}\qquad\Psi_+^s=\{\alpha_1+\alpha_2,\alpha_1+2\alpha_2,\alpha_2\}$$

and the pairs (β,γ) of orthogonal positive roots are

$$(\alpha_1,\alpha_1+2\alpha_2),\qquad(2\alpha_1+3\alpha_2,\alpha_2),\qquad(\alpha_1+3\alpha_2,\alpha_1+\alpha_2)$$

The corresponding graph reads


(α_1,α_2)
$(\alpha_1,\alpha_1+\alpha_2)$
$(\alpha_1+\alpha_2,\alpha_2)$
$(\alpha_1,\alpha_1+2\alpha_2)$
$(\alpha_1+2\alpha_2,\alpha_2)$
$(\alpha_1,\alpha_1+3\alpha_2)$
$(\alpha_1+\alpha_2,\alpha_1+2\alpha_2)$
$(\alpha_1+3\alpha_2,\alpha_2)$
$(\alpha_1,2\alpha_1+3\alpha_2)$
$(\alpha_1+\alpha_2,\alpha_1+3\alpha_2)$
$(2\alpha_1+3\alpha_2,\alpha_2)$
$(\alpha_1+\alpha_2,2\alpha_1+3\alpha_2)$
$(\alpha_1+2\alpha_2,\alpha_1+3\alpha_2)$
$(\alpha_1+2\alpha_2,2\alpha_1+3\alpha_2)$
$(\alpha_1+3\alpha_2,2\alpha_1+3\alpha_2)$

This yields an $\eta_{a,b}$ component of Ω_1^Ψ equal to

$$\begin{aligned}
&\eta_{\alpha_1,\alpha_2}\,[t_{\alpha_1},t_{\alpha_2}]\\
&+\eta_{\alpha_1+\alpha_2,\alpha_2}\,[t_{\alpha_1+\alpha_2}+t_{\alpha_1},t_{\alpha_2}]\\
&+\eta_{\alpha_1+2\alpha_2,\alpha_2}\,[t_{\alpha_1+2\alpha_2}+t_{\alpha_1+\alpha_2}+t_{\alpha_1},t_{\alpha_2}]\\
&+\eta_{\alpha_1,\alpha_1+3\alpha_2}\,[t_{\alpha_1},t_{\alpha_1+3\alpha_2}]\\
&+\eta_{\alpha_1+\alpha_2,\alpha_1+2\alpha_2}\,([t_{\alpha_1+\alpha_2},t_{\alpha_1+2\alpha_2}+t_{\alpha_2}]+[t_{\alpha_1},t_{\alpha_2}])
\end{aligned}$$

By (2.8), the first commutator is equal to $-[t_{\alpha_1},t_{\alpha_1+\alpha_2}]$. By (Φ) and $(\mathsf{A}_1\times\mathsf{A}_1\subset\mathsf{G}_2)$, the third commutator is equal to $-[t_{\alpha_1+3\alpha_2},t_{\alpha_2}]$. The second commutator is therefore equal to

$$-[t_{\alpha_1+3\alpha_2}+t_{\alpha_1+2\alpha_2},t_{\alpha_2}]=-[t_{\alpha_1+2\alpha_2},t_{\alpha_2}+t_{\alpha_1+3\alpha_2}]$$

where we used (2.8). By $(\mathsf{A}_2\subset G_2)$, the fourth commutator is equal to $-[t_{\alpha_1},t_{2\alpha_1+3\alpha_2}]$. Finally, the coefficient of $\eta_{\alpha_1+\alpha_2,\alpha_1+2\alpha_2}$ is equal to

$$-[t_{\alpha_1+\alpha_2},t_{\alpha_1}+t_{2\alpha_1+3\alpha_2}]+[t_{\alpha_1},t_{\alpha_2}]=[t_{2\alpha_1+3\alpha_2},t_{\alpha_1+\alpha_2}]$$

where we used (2.8). Using (2.17) shows that the right–hand side of (2.15) is equal to the $\eta_{a,b}$–component of Ω_1^Ψ. A similar use of relations (Φ), $(\mathsf{A}_2\subset G_2)$ and $(\mathsf{A}_1\times\mathsf{A}_1\subset\mathsf{G}_2)$ shows that the $\eta_a\wedge\eta_b$ component of Ω_1^Ψ is zero and therefore completes the proof ■

2.18.

Corollary. *Assume that the relations (tt) and (tτ) hold. Then, for any rank 2 root subsystem $\Psi\subset\Phi$, the curvature term Ω_1^Ψ is equal to*

$$\Omega_1^\Psi=-\sum_{\alpha\in\Psi_+}\sum_{\beta\in\Psi_+\cap w_\alpha\Psi_-}\eta_\alpha\wedge d\beta\,[t_\alpha,t_\beta]$$

where w_α is any element of the Weyl group of Ψ such that $w_\alpha^{-1}\alpha$ is simple in Ψ.

Proof. Lemma 2.17 and a simple case–by–case inspection show that Ω_1^Ψ does indeed have the above form for well–chosen elements w_α (specifically, w_α should be an element of shortest length such that $w_\alpha^{-1}\alpha$ is simple in Ψ). But Lemma 2.16 applied to Ψ implies that the expression

$$\sum_{\beta\in\Psi_+\cap w_\alpha\Phi_-}\eta_\alpha\wedge d\beta\,[t_\alpha,t_\beta]$$

is independent of the choice of w_α, hence the conclusion. ■

2.19. **Completion of the proof of (1) of Theorem 2.5.** Fix $\alpha\in\Phi_+$ and let $\mathcal{R}_2(\alpha)$ be the set of complete, rank 2 subsystems $\Psi\subset\Phi$ containing α as a non–simple root. For $\Psi\in\mathcal{R}_2(\alpha)$, denote the corresponding Weyl group by $W(\Psi)$.

For any $w\in W$, denote by $N(w)\subset\Phi_+$ the set

$$N(w)=\{\beta\in\Phi_+|\,w\beta\in\Phi_-\}$$

and define similarly $N_\Psi(w) \subset \Psi_+$ for any $w \in W(\Psi)$.

The following result, together with Lemma 2.16 and Corollary 2.18 show that $\Omega_1 + \Omega_2 = 0$ and therefore that the connection (2.2) is flat.

Proposition. *Let $w \in W$ be such that $w^{-1}\alpha$ is simple in Φ.*

1. *For any $\Psi \in \mathcal{R}_2(\alpha)$, the intersection $N(w^{-1}) \cap \Psi$ is non–empty.*
2. *The following holds*
$$N(w^{-1}) = \bigsqcup_{\Psi \in \mathcal{R}_2(\alpha)} N(w^{-1}) \cap \Psi$$
3. *For any $\Psi \in \mathcal{R}_2(\alpha)$, there exists a unique $w_\Psi \in W(\Psi)$ such that*
$$N(w^{-1}) \cap \Psi = N_\Psi(w_\Psi^{-1})$$

Proof. For any pair of non–proportional roots $\beta, \gamma \in \Phi$, let $\langle \beta, \gamma \rangle \subset \Phi$ be the complete, rank 2 subsystem generated by β and γ. We claim that the map $\beta \to \langle \alpha, \beta \rangle$ induces a bijection

$$\rho : N(w^{-1})/_\sim \longrightarrow \mathcal{R}_2(\alpha)$$

where $\sim$ is the equivalence relation defined by $\beta \sim \beta'$ if $\langle \alpha, \beta \rangle = \langle \alpha, \beta' \rangle$. This clearly proves (1) and (2) since $\rho^{-1}(\Psi) = \Psi \cap N(w^{-1})$.

To see this, we shall need some notation. For any complete subsystem $\Psi \subset \Phi$, let $\Psi^\perp = \bigcap_{\beta \in \Psi} \mathcal{H}_\beta \subset E$, where $\mathcal{H}_\beta = \mathrm{Ker}(\beta)$. Set $E_\Psi = E/\Psi^\perp$ so that Ψ may be regarded as a root system in E_Ψ^*, and let $\pi_\Psi : E \to E_\Psi$ be the corresponding projection. If $\beta \in N(w^{-1})$, the wall $\mathcal{H}_\beta$ separates the fundamental chamber $C \subset E$ of Φ and $C' = w(C)$. Thus, if $\Psi = \langle \alpha, \beta \rangle$, $\pi_\Psi(\mathcal{H}_\beta)$ separates the fundamental chamber $\pi_\Psi(C)$ of Ψ and $\pi_\Psi(C')$. It follows that $\pi_\Psi(C) \neq \pi_\Psi(C')$ so that α is not simple in Ψ since $\pi_\Psi(\mathcal{H}_\alpha)$ is a wall of $\pi_\Psi(C')$, whence $\langle \alpha, \beta \rangle \in \mathcal{R}_2(\alpha)$ and ρ is a well–defined embedding.

It is also surjective since if $\Psi \in \mathcal{R}_2(\alpha)$, there exists a $\beta \in \Psi_+$ such that $\pi_\Psi(\mathcal{H}_\beta)$ separates $\pi_\Psi(C)$ and $\pi_\Psi(C')$ so that $\mathcal{H}_\beta$ separates C and C' and therefore lies in $N(w^{-1})$.

Finally, for a given $\Psi \in \mathcal{R}_2(\alpha)$, the set $N(w^{-1}) \cap \Psi$ consists of those $\beta \in \Psi_+$ which separate C and C' and therefore $\pi_\Psi(C)$ and $\pi_\Psi(C')$. It is therefore equal to $\Psi_+ \cap w_\Psi \Psi_-$ where $w_\Psi \in W(\Psi)$ is the unique element such that $\pi_\Psi(C') = w_\Psi \pi_\Psi(C)$. ■

2.20. We now turn to part (2) of Theorem 2.5. We shall need the following

Lemma. *The relations (tt) imply that the following holds for any $\alpha \in \Phi_+$, $w \in W$ such that $w^{-1}\alpha$ is simple and $v \in \mathfrak{h}$ such that $\alpha(v) = 0$*

$$[t_\alpha, \sum_{\beta \in \Phi_+} \mathrm{sign}(w^{-1}\beta)\beta(v)t_\beta] = 0 \qquad (t^w t)$$

where $\mathrm{sign}(\gamma) = \pm 1$ *depending on whether* $\gamma \in \pm\Phi_+$.

Proof. Let $\mathcal{T}$ be the algebra generated by symbols t_α, $\alpha \in \Phi$ subject to the relations $t_{-\alpha} = t_\alpha$ and (tt). The Weyl group acts on $\mathcal{T}$ by $wt_\alpha = t_{w\alpha}$ and it is easy to check that the above relation holds for a triple (α, w, v) if, and only if, it holds for $(w^{-1}\alpha, 1, w^{-1}v)$. We may therefore assume that α is simple and that $w = 1$. Since the left–hand side of $(t^w t)$ may be written as

$$\sum_{\Psi} \sum_{\beta \in \Psi_+} [t_\alpha, \beta(v) t_\beta]$$

where Ψ ranges over the complete, rank 2 subsystems of Φ containing α, it is sufficient to prove $(t^w t)$ when Φ is of rank 2. In this case, it follows by a simple case–by–case verification. For example, if $\Phi_+ = \{\alpha_1, \alpha_2, \alpha_1 + \alpha_2, \alpha_1 + 2\alpha_2, \alpha_1 + 3\alpha_2, 2\alpha_1 + 3\alpha_2\}$ is of type G_2, we have

$$[t_{\alpha_2}, \sum_{\beta \in \Phi_+} \beta(\lambda_1^\vee) t_\beta] = [t_{\alpha_2}, \sum_{\beta \in \Phi_+} t_\beta] + [t_{\alpha_2}, t_{2\alpha_1 + 3\alpha_2}] = 0$$

by (Φ) and $(\mathsf{A}_1 \times \mathsf{A}_1 \subset \mathsf{G}_2)$, while

$$[t_{\alpha_1}, \sum_{\beta \in \Phi_+} \beta(\lambda_2^\vee) t_\beta] = [t_{\alpha_1}, \sum_{\beta \in \Phi_+} t_\beta] + [t_{\alpha_1}, t_{\alpha_1 + 2\alpha_2}] + 2[t_{\alpha_1}, t_{2\alpha_1 + 3\alpha_2} + t_{\alpha_1 + 3\alpha_2}]$$

which is equal to zero by (Φ), $(\mathsf{A}_1 \times \mathsf{A}_1 \subset \mathsf{G}_2)$ and $(\mathsf{A}_2 \subset G_2)$ ■

2.21. Recall from §2.3 that $\delta : \mathfrak{h} \to A$ is defined by

$$\delta(v) = \tau(v) - \frac{1}{2} \sum_{\alpha \in \Phi_+} \alpha(v) t_\alpha$$

The following proves part (2) of Theorem 2.5

Proposition. *Modulo the relations (tt), the relations $(t\tau)$ are equivalent to*

$$[t_\alpha, \delta(v)] = 0 \tag{$t\delta$}$$

for any $\alpha \in \Phi$ and $v \in \mathfrak{h}$ such that $\alpha(v) = 0$.

Proof. For any $w \in W$ (2.5), yields

$$\tau_w(v) = \tau(v) - \sum_{\alpha \in \Phi_+ \cap w\Phi_-} \alpha(v) t_\alpha = \delta(v) + \frac{1}{2} \sum_{\beta \in \Phi_+} \operatorname{sign}(w^{-1}\beta) \beta(v) t_\beta(v)$$

The result now follows from Lemma 2.20. ■

2.22. **Equivariance under W.** Assume now that the algebra A is acted upon by the Weyl group W of Φ.

Proposition.

(1) *The connection ∇ is W–equivariant if, and only if*

$$s_i(t_\alpha) = t_{s_i \alpha} \tag{2.18}$$
$$s_i(\tau(x)) - \tau(s_i x) = (\alpha_i, x) t_{\alpha_i} \tag{2.19}$$

for any $\alpha \in \Phi$, simple reflection $s_i \in W$ and $x \in \mathfrak{h}$.

(2) *Modulo* (2.18)*, the relation* (2.19) *is equivalent to the* W*–equivariance of the linear map* $\delta : \mathfrak{h} \to A$ *defined by* (2.6)*.*

Proof. (1) Since s_i permutes the set $\Phi_+ \setminus \{\alpha_i\}$ and, by (2.4)

$$\frac{1}{1-e^{-\alpha_i}} = \frac{1}{e^{\alpha_i}-1} + 1$$

we get

$$s_i^*\nabla = d - \sum_{\alpha\in\Phi_+} \frac{d\alpha}{e^\alpha - 1} s_i(t_\alpha) - d\alpha_i\, s_i(t_{\alpha_i}) - s_i(\tau(u^j))d(s_i u_j)$$

Requiring that $s_i^*\nabla = \nabla$ and taking residues along each subtorus $\{e^\alpha = 1\}$ yields (2.18). To proceed, note that

$$s_i(\tau(u^j))d(s_i u_j) = s_i(\tau(s_i u^j))d(u_j)$$

since $\tau(u^j)du_j$ is independent of the choice of the dual bases $\{u^j\}, \{u_j\}$. Thus, $s_i^*\nabla = \nabla$ reduces to

$$\tau(u^j)du_j = s_i(\tau(s_i u^j))du_j + t_{\alpha_i} d\alpha_i$$

which, upon being contracted along the tangent vector u^k yields (2.19) with $x = s_i u^j$.

(2) It is easy to check that the map $\overline{\tau}(x) = 1/2 \sum_{\alpha\in\Phi_+}(x,\alpha)t_\alpha$ satisfies (2.19). The result now follows since any two maps $\tau_i : \mathfrak{h} \to A$ satisfying (2.19) differ by a W–equivariant map. ■

2.23. **Flatness and equivariance.** The following is a direct corollary of Theorem 2.5 and Proposition 2.22

Theorem. *The trigonometric connection*

$$\nabla = d - \sum_{\alpha\in\Phi_+} \frac{d\alpha}{e^\alpha - 1}\, t_\alpha - d(u_i)\,\tau(u^i)$$

is flat and W*–equivariant if, and only if the following relations hold*

- *For any rank 2 root subsystem* $\Psi \subset \Phi$ *and* $\alpha \in \Psi$*,*
$$[t_\alpha, \sum_{\beta\in\Psi_+} t_\beta] = 0$$
- *For any* $u, v \in \mathfrak{h}$*,*
$$[\tau(u), \tau(v)] = 0$$
- *For any simple root* α_i *and* $u \in \operatorname{Ker}(\alpha_i)$*,*
$$[t_{\alpha_i}, \tau(u)] = 0$$
- *For any* $\alpha \in \Phi$ *and simple reflection* $s_i \in W$*,*
$$s_i(t_\alpha) = t_{s_i\alpha}$$
- *For any* $\alpha \in \Phi$ *and* $u \in \mathfrak{h}$*,*
$$s_i(\tau(u)) - \tau(s_i u) = (\alpha_i, u)t_{\alpha_i}$$

Remark. Theorem 2.23 was first proved by Cherednik in the special case when t_α is equal to the orthogonal reflection $s_\alpha \in W$ and shown to lead to the definition of the degenerate affine Hecke algebra of W [Ch1, Ch2].

3. The trigonometric Casimir connection

3.1. **The Yangian** $Y(\mathfrak{g})$ [Dr1]**.** Let $\mathfrak{g}$ be a finite–dimensional, simple Lie algebra over $\mathbb{C}$ and $(\cdot,\cdot)$ a non–degenerate, invariant bilinear form on $\mathfrak{g}$. Let $\hbar$ be a formal variable. The Yangian $Y(\mathfrak{g})$ is the associative algebra over $\mathbb{C}[\hbar]$ generated by elements $x, J(x)$, $x \in \mathfrak{g}$ subject to the relations

$$\lambda x + \mu y \quad (\text{in } Y(\mathfrak{g})) \;= \lambda x + \mu y \quad (\text{in } \mathfrak{g})$$

$$xy - yx = [x,y]$$

$$J(\lambda x + \mu y) = \lambda J(x) + \mu J(y)$$

$$[x, J(y)] = J([x,y])$$

$$[J(x), J([y,z])] + [J(z), J([x,y])] + [J(y), J([z,x])] = \\ \hbar^2([x,x_a],[[y,x_b],[z,x_c]])\{x^a, x^b, x^c\}$$

$$[[J(x), J(y)],[z,J(w)]] + [[J(z),J(w)],[x,J(y)]] = \\ \hbar^2([x,x_a],[[y,x_b],[[z,w],x_c]])\{x^a, x^b, J(x^c)\}$$

for any $x,y,z,w \in \mathfrak{g}$ and $\lambda,\mu \in \mathbb{C}$, where $\{x_a\}, \{x^a\}$ are dual bases of $\mathfrak{g}$ with respect to $(\cdot,\cdot)$ and

$$\{z_1,z_2,z_3\} = \frac{1}{24}\sum_{\sigma\in\mathfrak{S}_3} z_{\sigma(1)}z_{\sigma(2)}z_{\sigma(3)}$$

$Y(\mathfrak{g})$ is an $\mathbb{N}$–graded $\mathbb{C}[\hbar]$–algebra provided one sets $\deg(x) = 0$, $\deg(J(x)) = 1$ and $\deg(\hbar) = 1$.

3.2. **Drinfeld's new realisation of** $Y(\mathfrak{g})$ [Dr2]**.** Let $\mathfrak{h} \subset \mathfrak{g}$ be a Cartan subalgebra of $\mathfrak{g}$, $\Phi = \{\alpha\} \subset \mathfrak{h}^*$ the corresponding root system. Let $\{\alpha_i\}_{i\in\mathbf{I}}$ be a basis of simple roots of Φ and $a_{ij} = 2(\alpha_i,\alpha_j)/(\alpha_i,\alpha_i)$ the entries of the Cartan matrix $\mathbf{A}$ of $\mathfrak{g}$. Set $d_i = (\alpha_i,\alpha_i)/2$, so that $d_i a_{ij} = d_j a_{ji}$ for any $i,j \in \mathbf{I}$.

Let $\nu : \mathfrak{h} \to \mathfrak{h}^*$ be the isomorphism determined by the inner product $(\cdot,\cdot)$ and set $t_i = \nu^{-1}(\alpha_i) = d_i\alpha_i^\vee$. For any $i \in \mathbf{I}$, choose root vectors $x_i^\pm \in \mathfrak{g}_{\pm\alpha_i}$ such that $[x_i^+, x_i^-] = t_i$. Recall that $\mathfrak{g}$ has a (slightly non–standard) presentation in terms of the generators $t_i, x_i^\pm$ with relations

$$[t_i,t_j] = 0$$

$$[t_i, x_j^\pm] = \pm d_i a_{ij} x_j^\pm$$

$$[x_i^+, x_j^-] = \delta_{ij} t_i$$

$$\operatorname{ad}(x_i^\pm)^{1-a_{ij}} x_j^\pm = 0$$

The Yangian $Y(\mathbf{A})$ is the associative algebra over $\mathbb{C}[\hbar]$ with generators $X_{i,r}^{\pm}, T_{i,r}$, $i \in \mathbf{I}$, $r \in \mathbb{N}$ and relations

$$[T_{i,r}, T_{j,s}] = 0$$

$$[T_{i,0}, X_{j,s}^{\pm}] = \pm d_i a_{ij} X_{j,s}^{\pm}$$

$$[T_{i,r+1}, X_{j,s}^{\pm}] - [T_{i,r}, X_{j,s+1}^{\pm}] = \pm \frac{\hbar}{2} d_i a_{ij} (T_{i,r} X_{j,s}^{\pm} + X_{j,s}^{\pm} T_{i,r})$$

$$[X_{i,r}^{+}, X_{j,s}^{-}] = \delta_{ij} T_{i,r+s}$$

$$[X_{i,r+1}^{\pm}, X_{j,s}^{\pm}] - [X_{i,r}^{\pm}, X_{j,s+1}^{\pm}] = \pm \frac{\hbar}{2} d_i a_{ij} (X_{i,r}^{\pm} X_{j,s}^{\pm} + X_{j,s}^{\pm} X_{i,r}^{\pm})$$

$$\sum_{\pi} [X_{i,r_{\pi(1)}}^{\pm}, [X_{i,r_{\pi(2)}}^{\pm}, [\cdots, [X_{i,r_{\pi(m)}}^{\pm}, X_{j,s}^{\pm}] \cdots]] = 0$$

where $i \neq j$ in the last relation, $m = 1 - a_{ij}$, $r_1, \ldots, r_m \in \mathbb{N}$ is any sequence of non–negative integers, and the sum is over all permutations π of $\{1, \ldots, m\}$. $Y(\mathbf{A})$ is $\mathbb{N}$–graded by $\deg(T_{i,r}) = r = \deg(X_{i,r}^{\pm})$ and $\deg(\hbar) = 1$.

3.3. **Isomorphism between the two presentations** [Dr2]**.** Choose root vectors $x_\alpha \in \mathfrak{g}_\alpha$ for any $\alpha \in \Phi$ such that $(x_\alpha, x_{-\alpha}) = 1$ and let

$$\kappa_\alpha = x_\alpha x_{-\alpha} + x_{-\alpha} x_\alpha \tag{3.1}$$

be the truncated Casimir operator of the $\mathfrak{sl}_2$–subalgebra of $\mathfrak{g}$ corresponding to α. Then, the assignment

$$\varphi(t_i) = T_{i,0}, \quad \varphi(x_i^{\pm}) = X_{i,0}^{\pm}$$

$$\varphi(J(t_i)) = T_{i,1} + \hbar \varphi(v_i)$$

$$\varphi(J(x_i^{\pm})) = X_i^{\pm} + \hbar \varphi(w_i^{\pm})$$

extends to an isomorphism $\varphi : Y(\mathfrak{g}) \to Y(\mathbf{A})$, where

$$v_i = \frac{1}{4} \sum_{\beta \in \Phi_+} (\alpha_i, \beta) \kappa_\alpha - t_i^2/2$$

$$w_i^{\pm} = \pm \frac{1}{4} \sum_{\beta \in \Phi_+} \left([x_i^{\pm}, x_{\pm\beta}] x_{\mp\beta} + x_{\mp\beta} [x_i^{\pm}, x_{\pm\beta}] \right) - \frac{1}{4} (x_i^{\pm} t_i + t_i x_i^{\pm})$$

3.4. **The W–equivariant embedding $\delta_{a,b} : \mathfrak{h} \to Y(\mathfrak{g})$.** The Yangian $Y(\mathfrak{g})$ is acted upon by the Lie algebra spanned by the elements $x \in \mathfrak{g}$ and is an integrable $\mathfrak{g}$–module under this action. In particular, the zero–weight subalgebra $Y(\mathfrak{g})^{\mathfrak{h}}$ is acted upon by the Weyl group W of $\mathfrak{g}$. Moreover, for any $a, b \in \mathbb{C}$, the linear map

$$\delta_{a,b} : \mathfrak{h} \to Y(\mathfrak{g})^{\mathfrak{h}}, \qquad \delta_{a,b}(t) = at + bJ(t) \tag{3.2}$$

is W–equivariant.

In terms of the new realisation of $Y(\mathfrak{g})$, the map $\delta_{a,b}$ becomes

$$\widetilde{\delta}_{a,b} = \varphi \circ \delta_{a,b} : \mathfrak{h} \to Y(\mathbf{A})^{\mathfrak{h}}$$

$$\widetilde{\delta}_{a,b}(t) = at + b\left(T(t)_1 + \frac{\hbar}{4} \sum_{\beta\in\Phi_+} \beta(t)\kappa_\beta - \frac{\hbar}{2}\sum_i (t, t^i) t_i^2 \right) \tag{3.3}$$

where $\{t^i = \lambda_i^\vee\}$ is the basis of $\mathfrak{h}$ dual to $\{t_i\}$ given by the fundamental coweights, $T(-)_1 : \mathfrak{h} \to Y(\mathbf{A})$ is the embedding $t \to (t, t^i)T_{i,1}$ and, deviating slightly from the notation of §3.2, we identify $\mathfrak{g} \subset Y(\mathfrak{g})$ with the Lie subalgebra of $Y(\mathbf{A})$ spanned by $T_{i,0}, X^{\pm}_{i,0}$.

3.5. **The linear map $\tau_{a,b} : \mathfrak{h} \to Y(\mathfrak{g})$.** Let $\epsilon \in \mathbb{C}$. For any root α, set

$$K_\alpha = \kappa_\alpha + \epsilon q_\alpha \qquad \text{where} \qquad q_\alpha = \frac{\nu^{-1}(\alpha)^2}{(\alpha,\alpha)} \tag{3.4}$$

and κ_α is the truncated Casimir given by (3.1). For any $a, b \in \mathbb{C}$, define a map $\tau_{a,b} : \mathfrak{h} \to Y(\mathfrak{g})^{\mathfrak{h}}$ by

$$\tau_{a,b}(t) = \frac{\hbar}{2} \sum_{\alpha\in\Phi_+} (t,\alpha) K_\alpha + \delta_{a,b}(t)$$

where $\delta_{a,b}$ is given by (3.2).

Proposition.

(1) *The elements* K_α *satisfy*

$$K_{-\alpha} = K_\alpha \qquad \text{and} \qquad w(K_\alpha) = K_{w\alpha}$$

for any $w \in W$.

(2) *The following holds for any* $t \in \mathfrak{h}$

$$s_i(\tau_{a,b}(t)) - \tau_{a,b}(s_i t) = \hbar(\alpha_i, t) K_{\alpha_i}$$

(3) *If* $b = -2$*, then* $\widetilde{\tau}_{a,b} = \varphi \circ \tau_{a,b} : \mathfrak{h} \to Y(\mathbf{A})^{\mathfrak{h}}$ *is given by*

$$\widetilde{\tau}_{a,-2}(t) = at - 2\left(T(t)_1 - \frac{\hbar}{2}\sum_i (t, t^i) t_i^2 \right) + \epsilon \frac{\hbar}{2} \sum_{\alpha\in\Phi_+} \alpha(t) q_\alpha$$

and satisfies in addition

$$[\widetilde{\tau}_{a,-2}(t), \widetilde{\tau}_{a,-2}(t')] = 0$$

for any $t, t' \in \mathfrak{h}$.

(4) *For any* $\alpha \in \Phi_+$ *and* $t \in \mathfrak{h}$ *such that* $\alpha(t) = 0$

$$[K_\alpha, \delta_{a,b}(t)] = 0$$

Proof. (1) is obvious. (2) follows from the second part of Proposition 2.22, (1) and the W–equivariance of $\delta_{a,b}$. For (3), we have by (3.3)

$$\begin{aligned}\widetilde{\tau}_{a,b}(t) &= \frac{\hbar}{2}\sum_{\alpha\in\Phi_+}\alpha(t)K_\alpha + at + b\left(T(t)_1 + \frac{\hbar}{4}\sum_{\beta\in\Phi_+}\beta(t)\kappa_\beta - \frac{\hbar}{2}\sum_i (t,t^i)t_i^2\right)\\ &= at + \frac{\hbar}{2}\sum_{\alpha\in\Phi_+}\alpha(t)(K_\alpha + \frac{b}{2}\kappa_\alpha) + b(T(t)_1 - \frac{\hbar}{2}\sum_i (t,t^i)t_i^2)\end{aligned}$$

which, for $b=-2$ reduces to the claimed expression. The commutativity of $\widetilde{\tau}_{a,-2}(t)$ and $\widetilde{\tau}_{a,-2}(t')$ then follows from that of of the $T_{i,1}$. (4) follows easily from the defining relations of $Y(\mathfrak{g})$. ■

3.6. For simplicity, we henceforth set $\epsilon = 0 = a$ and $b=-2$ in equations (3.2) and (3.4), although Theorem 3.8 below is true for any values of a,ϵ. Thus, $K_\alpha = \kappa_\alpha$, $\delta = \delta_{0,-2} : \mathfrak{h}\to Y(\mathfrak{g})$ is given by $\delta(t) = -2J(t)$ and the corresponding map $\widetilde{\tau} = \widetilde{\tau}_{0,-2} : \mathfrak{h} \to Y(\mathbf{A})^{\mathfrak{h}}$ by

$$\widetilde{\tau}(t) = -2T(t)_1 + \hbar(t,t^i)t_i^2$$

3.7. **The trigonometric Casimir connection of $\mathfrak{g}$.** Let $H_{\mathrm{reg}}\subset H$ be given by (2.1) and $\mathcal{Y}^{\mathfrak{h}}$ the trivial bundle over H_{reg} with fibre $Y(\mathfrak{g})^{\mathfrak{h}} \cong Y(\mathbf{A})^{\mathfrak{h}}$.

Definition. The *trigonometric Casimir connection* of $\mathfrak{g}$ is the connection ∇ on $\mathcal{Y}^{\mathfrak{h}}$ given by either of the following forms

$$\nabla = d - \frac{\hbar}{2}\sum_{\alpha\in\Phi_+}\frac{e^\alpha+1}{e^\alpha-1}d\alpha\,\kappa_\alpha + 2du_i\,J(u^i) \tag{3.5}$$

$$= d - \hbar\sum_{\alpha\in\Phi_+}\frac{d\alpha}{e^\alpha-1}\,\kappa_\alpha + 2du_i\left(T(u^i)_1 - \frac{\hbar}{2}(u^i,t^j)t_j^2\right) \tag{3.6}$$

whose equality follows from §2.3 and §3.6.

3.8.

Theorem. *The trigonometric Casimir connection is flat and W–equivariant.*

Proof. By Theorem 2.5, Proposition 2.22 and Proposition 3.5, we need only check that the elements $t_\alpha = \kappa_\alpha$ satisfy the relations (tt). This follows as in [MTL, Thm. 2.3] and [TL1, Thm 2.2]. Specifically, if $\Psi\subset\Phi$ is a rank 2 root subsystem, the sum $\sum_{\alpha\in\Phi_+}\kappa_\alpha$ is, up to Cartan terms, the Casimir operator for the rank 2 subalgebra $\mathfrak{g}_\Psi\subset\mathfrak{g}$ determined by Ψ and therefore commutes with each summand κ_α, $\alpha\in\Phi_+$. ■

Remark. Since the relations of Theorem 2.5 and Proposition 2.22 are homogeneous, Theorem 3.8 proves in fact the flatness and W–equivariance of the one–parameter family of connections

$$\begin{aligned}\nabla &= d - \lambda^{-1}\left(\frac{\hbar}{2}\sum_{\alpha\in\Phi_+}\frac{e^\alpha+1}{e^\alpha-1}d\alpha\,\kappa_\alpha - 2du_i\,J(u^i)\right)\\ &= d - \lambda^{-1}\left(\hbar\sum_{\alpha\in\Phi_+}\frac{d\alpha}{e^\alpha-1}\,\kappa_\alpha - 2du_i\left(T(u^i)_1 - \frac{\hbar}{2}(u^i,t^j)t_j^2\right)\right)\end{aligned}$$

where λ varies in $\mathbb{C}^\times$.

4. The monodromy conjecture

We show in this section that the monodromy of the trigonometric Casimir connection ∇ gives rise to representations of the affine braid group $\widehat{B}$ corresponding to $\mathfrak{g}$ and give a conjectural description of it in terms of the quantum Weyl group operators of the quantum loop algebra $U_\hbar(L\mathfrak{g})$.

4.1. **Monodromy representation.** Since H is of simply–connected type, the Weyl group W acts freely on H_{reg} and the fundamental group of the quotient H_{reg}/W is isomorphic to the affine braid group $\widehat{B}$ [NVD, vdL].

Let V be a finite–dimensional $Y(\mathfrak{g})$–module and $\mathbb{V}$ the holomorphically trivial vector bundle over H_{reg} with fibre V. The connection ∇ induces a flat connection on $\mathbb{V}$. To push it down to the quotient by W we use the 'up and down' trick of [MTL, p. 224] to circumvent the fact that W does not in general act on V. To this end, we shall need a few basic results about Tits extensions of (affine) Weyl groups which are gathered in §8.

Specifically, since V is an integrable $\mathfrak{g}$–module, the triple exponentials

$$\exp(e_{\alpha_i})\exp(-f_{\alpha_i})\exp(e_{\alpha_i})$$

defined by a choice of simple root vectors $e_{\alpha_i}\in\mathfrak{g}_{\alpha_i}, f_{\alpha_i}\in\mathfrak{g}_{-\alpha_i}$ are well–defined elements of $GL(V)$. They give rise to an action on V of an extension of W by the sign group $\mathbb{Z}_2^{\dim\mathfrak{h}}$ called the *Tits extension* $\widetilde{W}$ of W (Definition 8.2 and Prop. 8.3). By Theorem 8.10, $\widetilde{W}$ is a quotient of the affine braid group $\widehat{B}$ which may therefore be made to act on V. It is then easy to check that the pull–back of the flat vector bundle $(\mathbb{V},\nabla)$ to the universal cover of H_{reg} is equivariant under $\widehat{B}$ acting by deck transformations on the base and through the $\widetilde{W}$–action on the fibres.

4.2. Let $L\mathfrak{g}=\mathfrak{g}[t,t^{-1}]$ be the loop algebra of $\mathfrak{g}$ and $U_\hbar(L\mathfrak{g})$ the corresponding quantum loop algebra, viewed as a topological Hopf algebra over the ring of formal power series $\mathbb{C}[\![\hbar]\!]$. Thus, $U_\hbar(L\mathfrak{g})$ has Chevalley generators E_i, F_i, where i ranges over the set $\widehat{\mathbf{I}}=\mathbf{I}\sqcup\{0\}$ of nodes of the affine Dynkin diagram of $\mathfrak{g}$ and a Cartan subalgebra isomorphic to $\mathfrak{h}$ and spanned by H_i,

$i \in \mathbf{I}$ and $H_0 = -H_\theta = -\sum_{i\in\mathbf{I}} a_i H_i$, where $\theta \in \mathfrak{h}^*$ is the highest root and the integers a_i are given by $\theta^\vee = \sum_i a_i \alpha_i^\vee$.

4.3. By a *finite–dimensional representation* of $U_\hbar(L\mathfrak{g})$ we shall mean a module $\mathcal{V}$ which is topologically free and finitely–generated over $\mathbb{C}[\![\hbar]\!]$. Such a $\mathcal{V}$ is integrable and therefore endowed with a quantum Weyl group action of the affine braid group $\widehat{B}$ [Lu, KR, So]. This action is given by letting the generator corresponding to $i \in \widehat{\mathbf{I}}$ act by

$$\overline{S}_i^\hbar\, v = \sum_{\substack{a,b,c\in\mathbb{Z}:\\ a-b+c=-\lambda(\alpha_i^\vee)}} (-1)^b q_i^{b-ac} E_i^{(a)} F_i^{(b)} E_i^{(c)} v$$

where $v \in \mathcal{V}$ if of weight $\lambda \in \mathfrak{h}^*$ and $X_i^{(a)}$ is the divided power $X^a/[a]_i!$ with

$$q = e^\hbar, \qquad q_i = q^{(\alpha_i,\alpha_i)/2}$$

$$[n]_i = \frac{q_i^n - q_i^{-n}}{q_i - q_i^{-1}} \qquad \text{and} \qquad [n]_i! = [n]_i[n-1]_i \cdots [1]_i$$

4.4. **Monodromy conjecture.** It is known that the Yangian $Y(\mathfrak{g})$ and the quantum loop algebra $U_\hbar(L\mathfrak{g})$ have the same finite–dimensional representation theory (see [Va] and [GTL1]). By analogy with the quantum Weyl group description of the monodromy of the (rational) Casimir connection of $\mathfrak{g}$ conjectured by De Concini (unpublished) and independently in [TL1, TL2], and proved in [TL1, TL3], we make the following

Conjecture. *The monodromy of the trigonometric Casimir connection is equivalent to the quantum Weyl group action of the affine braid group* $\widehat{B}$ *on finite–dimensional* $U_\hbar(L\mathfrak{g})$*–modules.*

We will return to this conjecture in forthcoming work in collaboration with S. Gautam [GTL2].

5. The trigonometric Casimir connection of $\mathfrak{gl}_n$

We consider in this section the Yangian $Y(\mathfrak{gl}_n)$ of the Lie algebra $\mathfrak{gl}_n$. The latter does not possess a presentation of the form given in §3.1 but may be defined via a ternary, or *RTT* presentation. By exploiting the interplay between the latter and its loop presentation, we construct a flat, trigonometric connection with values in $Y(\mathfrak{gl}_n)$. We then relate it to the corresponding connection for $\mathfrak{sl}_n$ and show that, when it is taken with values in a tensor product of evaluation modules, it coincides with the trigonometric dynamical equations [TV].

5.1. **The RTT presentation of $Y(\mathfrak{gl}_n)$.** The Yangian $Y(\mathfrak{gl}_n)$ is the unital, associative algebra over $\mathbb{C}$ generated by elements $t_{ij}^{(r)}$, $1 \leq i,j \leq n$, $r \geq 1$, subject to the relations [3]

$$[t_{ij}^{(r+1)}, t_{kl}^{(s)}] - [t_{ij}^{(r)}, t_{kl}^{(s+1)}] = t_{kj}^{(r)} t_{il}^{(s)} - t_{kj}^{(s)} t_{il}^{(r)} \tag{5.1}$$

where $r, s \in \mathbb{N}$ and $t_{ij}^{(0)} = \delta_{ij}$.

Let $V = \mathbb{C}^n$ with standard basis $e_1, \ldots, e_n$ and let $E_{ij} e_k = \delta_{jk} e_i$ be the corresponding basis of elementary matrices of $\mathfrak{gl}_n$. The map $\imath : E_{ij} \to t_{ij}^{(1)}$ defines an embedding of $\mathfrak{gl}_n$ into $Y(\mathfrak{gl}_n)$ and we will often identify $\mathfrak{gl}_n$ with its image under $\imath$. Moreover, for every $s \geq 1$, the subspace spanned by the elements $t_{ij}^{(s)}$ transforms like the adjoint representation under the commutator action of $\mathfrak{gl}_n$.

The above relations may be more compactly rewritten as follows. For any $r \geq 0$, let $T^{(r)}$ be the $n \times n$ matrix with values in $Y(\mathfrak{gl}_n)$ given by

$$T^{(r)} = \sum_{1 \leq i,j \leq n} E_{ij}^V \otimes t_{ij}^{(r)}$$

where E_{ij}^V are again elementary matrices and the superscript V is used to stress the fact that they should be thought of as elements of the algebra $\mathrm{End}(V)$ rather than the underlying Lie algebra $\mathfrak{gl}_n$.

Let u be a formal variable and set

$$T = \sum_{r \geq 0} T^{(r)} u^{-r} \in \mathrm{End}(V) \otimes Y(\mathfrak{gl}_n)[\![u^{-1}]\!]$$

Finally, let

$$R(u) = 1 - P u^{-1} \in \mathrm{End}(V \otimes V)[\![u^{-1}]\!]$$

be Yang's R–matrix, where $P \in \mathrm{End}(V \otimes V)$ acts as the permutation of the two tensor factors. Then the relations (5.1) are equivalent to

$$R(u-v) T_1(u) T_2(v) = T_2(v) T_1(u) R(u-v)$$

where $T_1(u), T_2(u) \in \mathrm{End}(V \otimes V) \otimes Y(\mathfrak{gl}_n)[\![u^{-1}]\!]$ are given by

$$T_1(u) = \sum_{i,j,r} E_{ij}^V \otimes 1 \otimes t_{ij}^{(r)} u^{-r} \qquad \text{and} \qquad T_2(v) = \sum_{i,j,r} 1 \otimes E_{ij}^V \otimes t_{ij}^{(r)} v^{-r}$$

5.2. **The loop presentation of $Y(\mathfrak{gl}_n)$.** Let $E(u), H(u), F(u)$ be the factors of the Gauss decomposition of $T(u)$. Specifically,

$$F(u) = 1 + \sum_{i>j} E_{ij}^V \otimes f_{ij}(u) \qquad E(u) = 1 + \sum_{i<j} E_{ij}^V \otimes e_{ij}(u)$$

$$H(u) = \sum_i E_{ii}^V \otimes h_i(u),$$

---

[3] we follow here the conventions of [Mo] and [NO].

are, respectively, the unique lower unipotent, upper unipotent and diagonal matrices with coefficients in $Y(\mathfrak{gl}_n)[\![u^{-1}]\!]$ such that

$$T(u) = F(u)H(u)E(u) \tag{5.2}$$

Noting that $H(u), E(u), F(u) = 1 \mod u^{-1}$, write

$$h_i(u) = 1 + \sum_{r\geq 1} h_i^{(r)} u^{-r}, \quad f_{ij}(u) = \sum_{r\geq 1} f_{ij}^{(r)} u^{-r}, \quad e_{ij}(u) = \sum_{r\geq 1} e_{ij}^{(r)} u^{-r}$$

The coefficients of $e_{ii+1}(u), f_{ii+1}(u)$ and $h_i(u)$ give another system of generators of $Y(\mathfrak{gl}_n)$. Moreover, The elements $h_i(u)$ commute and their coefficients generate a maximal commutative subalgebra of $Y(\mathfrak{gl}_n)$ called the *Gelfand–Zetlin subalgebra* H_n.

5.3. The Gauss decomposition (5.2) yields in particular

$$t_{ij}^{(1)} = \begin{cases} e_{ij}^{(1)} & \text{if } i < j \\ h_i^{(1)} & \text{if } i = j \\ f_{ij}^{(1)} & \text{if } i > j \end{cases}$$

which we will use to identify the copies of $\mathfrak{gl}_n$ inside each presentation. Moreover,

$$\begin{aligned} t_{ii}^{(2)} &= h_i^{(2)} + \sum_{j<i} E_{ij}E_{ji} \\ &= h_i^{(2)} + \frac{1}{2}\sum_{j<i} \left(\kappa_{\theta_j-\theta_i} - (E_{jj} - E_{ii})\right) \end{aligned} \tag{5.3}$$

where θ_a is the linear form given by $\theta_a(E_{bb}) = \delta_{ab}$ and $\kappa_{\theta_a-\theta_b} = E_{ab}E_{ba} + E_{ba}E_{ab}$ is the truncated Casimir operator corresponding to the root $\theta_a - \theta_b$.

5.4. Define the elements $D_i \in H_n$ by

$$D_i = 2h_i^{(2)} - \sum_{j<i}(E_{jj} - E_{ii}) - E_{ii}^2 \tag{5.4}$$

$$= 2t_{ii}^{(2)} - \sum_{j<i} \kappa_{\theta_j-\theta_i} - E_{ii}^2 \tag{5.5}$$

The symmetric group $\mathfrak{S}_n$ acts by algebra automorphisms on $Y(\mathfrak{gl}_n)$ by

$$\sigma(t_{ij}^{(r)}) = t_{\sigma(i)\sigma(j)}^{(r)} \tag{5.6}$$

Lemma. *The following holds*

1. $[D_i, D_j] = 0$.
2. $(i\,i+1)D_j = D_j$ *if* $j \notin \{i, i+1\}$.
3. $(i\,i+1)D_i - D_{i+1} = \kappa_{\theta_i-\theta_{i+1}}$.

Proof. (1) follows from (5.4) and the fact that the $h_i^{(r)}$ commute. (2) and (3) follows from (5.5). ■

5.5. The following is a direct consequence of (5.5) and (5.3)

Lemma. *The element* $\mathbf{D} = D_1 + \cdots + D_n$ *is given by*

$$\begin{aligned}\mathbf{D} &= 2\sum_i t_{ii}^{(2)} - C_{\mathfrak{gl}_n}\\ &= 2\sum_i h_i^{(2)} - 2\rho^\vee - \sum_i E_{ii}^2\end{aligned}$$

where

$$C_{\mathfrak{gl}_n} = \sum_{i<j}\kappa_{\theta_i-\theta_j} + \sum_i E_{ii}^2 \qquad \textit{and} \qquad 2\rho^\vee = \sum_{i<j}(E_{ii}-E_{jj})$$

are the Casimir operator and sum of the positive coroots of $\mathfrak{gl}_n$*.*

5.6. **The trigonometric Casimir connection of $\mathfrak{gl}_n$.** Let D_i be given by (5.5) and define elements $\Delta_i \in H_n$ by (cf. §2.3)

$$\begin{aligned}\Delta_i &= D_i - \frac{1}{2}\sum_{a<b}(\theta_a-\theta_b)(E_{ii})\,\kappa_{\theta_a-\theta_b}\\ &= 2t_{ii}^{(2)} - \frac{1}{2}\sum_{j\neq i}\kappa_{\theta_i-\theta_j} - E_{ii}^2\end{aligned}$$

Let $H \subset GL_n$ be the maximal torus consisting of diagonal matrices, H_{reg} its set of regular elements and $\mathcal{Y}(\mathfrak{gl}_n)$ the trivial $Y(\mathfrak{gl}_n)$–bundle over H_{reg}.

Definition. The *trigonometric Casimir connection* of $\mathfrak{gl}_n$ is the connection on $\mathcal{Y}(\mathfrak{gl}_n)$ given by either of the following forms

$$\nabla = d - \sum_{i<j}\frac{d(\theta_i-\theta_j)}{e^{\theta_i-\theta_j}-1}\,\kappa_{\theta_i-\theta_j} - \sum_{i=1}^n d\theta_i\,D_i$$

$$= d - \frac{1}{2}\sum_{i<j}\frac{e^{\theta_i-\theta_j}+1}{e^{\theta_i-\theta_j}-1}d(\theta_i-\theta_j)\,\kappa_{\theta_i-\theta_j} - \sum_{i=1}^n d\theta_i\,\Delta_i$$

5.7. Let the symmetric group $\mathfrak{S}_n$ act on the vector bundle $\mathcal{Y}(\mathfrak{gl}_n)$ by permutations of the base and automorphisms (5.6) of the fibre.

Theorem. *The trigonometric Casimir connection of* $\mathfrak{gl}_n$ *is flat and equivariant under* $\mathfrak{S}_n$*.*

Proof. Let $\overline{H}, U \subset H$ be the subtori consisting respectively of diagonal matrices of determinant 1 and multiples of the identity, and let $\overline{\mathfrak{h}}, \mathfrak{u} = \mathbb{C}\mathbf{1_n}$ and $\mathfrak{h}$ be their Lie algebras, where $\mathbf{1_n} = \sum_{i=1}^n E_{ii}$. Thus

$$\mathfrak{h} = \overline{\mathfrak{h}} \oplus \mathbb{C}\mathbf{1_n} \qquad \text{and} \qquad \mathfrak{h}^* \cong \overline{\mathfrak{h}}^* \oplus \mathbb{C}\,\mathrm{tr}$$

where $\mathrm{tr} : \mathfrak{h} \to \mathbb{C}$ is the trace. Clearly, $H \cong \overline{H} \times U$ and the connection ∇ decomposes as the product of the following $Y(\mathfrak{gl}_n)$–valued connections on

$\overline{H}_{\text{reg}}$ and U respectively

$$\overline{\nabla} = d - \sum_{i<j} \frac{d(\theta_i - \theta_j)}{e^{\theta_i - \theta_j} - 1}\, \kappa_{\theta_i - \theta_j} - du_a\, D(u^a) \tag{5.7}$$

$$\nabla^U = d - \frac{1}{n} d \operatorname{tr} \mathbf{D} \tag{5.8}$$

where $D : \mathfrak{h} \to Y(\mathfrak{gl}_n)$ is given by $D(E_{ii}) = D_i$ and $\{u_a\}, \{u^a\}$ are dual bases of $\overline{\mathfrak{h}}^*$ and $\overline{\mathfrak{h}}$ respectively. ∇^U is clearly flat and equivariant under the action of $\mathfrak{S}_n$ since the latter acts trivially on U and, by Lemma 5.5 on $\mathbf{D}$. Since the latter commutes with the coefficients of $\overline{\nabla}$, the flatness and equivariance of ∇ reduces to that of $\overline{\nabla}$ which, in turn is determined by Theorem 2.5 and Proposition 2.22. The relations (tt) have already been checked in the proof of the flatness of the trigonometric Casimir connection for $\mathfrak{sl}_n$ in Theorem 3.8. The relations $(\tau\tau)$ and the equivariance relations (2.19) follow from Lemma 5.4. There remains to check that, for any $i = 1, \ldots, n-1$ and $u \in \operatorname{Ker}(\theta_i - \theta_{i+1})$, $[\kappa_{\theta_i - \theta_{i+1}}, D(u)] = 0$. This reduces to checking that $[\kappa_{\theta_i - \theta_{i+1}}, D_j] = 0$ for $j \notin \{i, i+1\}$ and that $[\kappa_{\theta_i - \theta_{i+1}}, D_i + D_{i+1}] = 0$ which follows easily from (5.5). ■

5.8. **Rational form.** It is well known that trigonometric connections of type GL_n may be put into a rational form, thus expressing them as KZ type connections on $n+1$ points, with one frozen to 0. We carry this step below for the connection 5.6.

Let $z_i = e^{\theta_i}$, $i = 1, \ldots, n$ be the standard coordinates on the torus $H \cong (\mathbb{C}^*)^n$ of GL_n.[4] Since $d\theta_i = dz_i / z_i$ and

$$\frac{d(\theta_i - \theta_j)}{e^{\theta_i - \theta_j} - 1} = \frac{d(z_i - z_j)}{z_i - z_j} - \frac{dz_i}{z_i}$$

the connection 5.6 is equal to

$$\nabla = d - \sum_{i<j} \frac{d(z_i - z_j)}{z_i - z_j}\, \kappa_{\theta_i - \theta_j} - \sum_i \frac{dz_i}{z_i}\, \widetilde{D}_i$$

where

$$\widetilde{D_i} = D_i - \sum_{j>i} \kappa_{\theta_i - \theta_j} = 2t^{(2)}_{ii} - \sum_{j \neq i} \kappa_{\theta_i - \theta_j} - E_{ii}^2$$

5.9. **Relation to $Y(\mathfrak{sl}_n)$.** Following Olshanski and Drinfeld, we realise the Yangian $Y(\mathfrak{sl}_n)$ as a Hopf subalgebra of $Y(\mathfrak{gl}_n)$ as follows (see [Mo, §1.8]). Let $\mathbb{A} = 1 + u^{-1}\mathbb{C}[\![u^{-1}]\!]$ be the abelian group of formal power series in u^{-1} with constant term 1. $\mathbb{A} \ni f$ acts on $Y(\mathfrak{gl}_n)$ by Hopf algebra automorphisms given by

$$T(u) \to f(u) T(u)$$

The Hopf subalgebra $Y(\mathfrak{gl}_n)^{\mathbb{A}} \subset Y(\mathfrak{gl}_n)$ of elements fixed by $\mathbb{A}$ is isomorphic to $Y(\mathfrak{sl}_n)$.

---

[4] note that these differ from the coordinates $z_i = e^{\alpha_i} = e^{\theta_i - \theta_{i+1}}$ used in §2.12.

5.10. The generators $T_{i,r}$ of the presentation of $Y(\mathfrak{sl}_n)$ described in §3.2 may be obtained within the RTT realisation of $Y(\mathfrak{gl}_n)$ as follows [BK, Rk. 5.12]. Consider their generating function $T_i(u) = 1+\sum_{r\geq 0} T_{i,r}u^{-r-1}$. Then,

$$T_i(u) = h_i(u - \frac{i-1}{2})^{-1} \cdot h_{i+1}(u - \frac{i-1}{2})$$

To spell this out, consider a formal power series $a(u) = 1 + a_1 u^{-1} + a_2 u^{-2} + \cdots$. Then, for any $\lambda \in \mathbb{C}$ one has

$$\begin{aligned} a(u-\lambda) &= 1 + a_1 u^{-1}(1 - \frac{\lambda}{u})^{-1} + a_2 u^{-2}(1 - \frac{\lambda}{u})^{-2} + \cdots \\ &= 1 + a_1 u^{-1} + (a_2 + \lambda a_1)u^{-2} + \cdots \end{aligned}$$

and therefore

$$a(u-\lambda)^{-1} = 1 - a_1 u^{-1} - (a_2 + \lambda a_1 - a_1^2)u^{-2} + \cdots$$

It follows that $T_{i,0} = -(E_{ii} - E_{i+1,i+1})$ and

$$T_{i,1} = -(h_i^{(2)} - h_{i+1}^{(2)}) - \frac{i-1}{2}(E_{ii} - E_{i+1,i+1}) + E_{ii}^2 - E_{ii}E_{i+1,i+1} \qquad (5.9)$$

5.11. Let $\overline{H} \subset H$ be the torus of SL_n consisting of diagonal matrices with determinant 1.

Proposition. *The restriction of the trigonometric Casimir connection of* $\mathfrak{gl}_n$ *to* $\overline{H}_{\text{reg}}$ *takes values in* $Y(\mathfrak{sl}_n)$ *and is equal to the sum of the trigonometric Casimir connection of* $\mathfrak{sl}_n$ *with the* $\overline{\mathfrak{h}} \subset Y(\mathfrak{sl}_n)$*–valued, closed one–form*

$$-\sum_{i=1}^{n} d\lambda_i \, (E_{ii} - E_{i+1,i+1})$$

where $\{\lambda_i\}$ *are the fundamental weights of* $\mathfrak{sl}_n$.

PROOF. The restriction of the trigonometric Casimir connecton of $\mathfrak{gl}_n$ to $\overline{H}_{\text{reg}}$ is given by (5.7), namely

$$\overline{\nabla} = d - \sum_{i<j} \frac{d(\theta_i - \theta_j)}{e^{\theta_i - \theta_j} - 1} \, \kappa_{\theta_i - \theta_j} - du_i \, D(u^i)$$

where $D : \mathfrak{h} \to Y(\mathfrak{gl}_n)$ is given by $D(E_{ii}) = D_i$ and $\{u_i\}, \{u^i\}$ are dual bases of $\overline{\mathfrak{h}}^*$ and $\overline{\mathfrak{h}}$ respectively. Choosing $u_i = \lambda_i$ so that $u^i = E_{ii} - E_{i+1,i+1}$, $i = 1, \ldots, n-1$ and comparing with the form (3.6) we need to show that

$$D_i - D_{i+1} = -2T_{i,1} + (E_{ii} - E_{i+1,i+1})^2 + (E_{ii} - E_{i+1,i+1})$$

By (5.4), the left–hand side is equal to

$$2(h_i^{(2)} - h_{i+1}^{(2)}) + i \, (E_{ii} - E_{i+1,i+1}) - E_{ii}^2 + E_{i+1,i+1}^2$$

and the result follows from (5.9) ■

5.12. **Evaluation homomorphism.** The Yangian $Y(\mathfrak{gl}_n)$ possesses an evaluation homomorphism $\mathrm{ev}: Y(\mathfrak{gl}_n)\rightarrow U\mathfrak{gl}_n$ defined by

$$\mathrm{ev}(t_{ij}(u)) = \delta_{ij} + E_{ij}u^{-1}$$

where $t_{ij}(u) = \sum_{r\geq 0} t_{ij}^{(r)}u^{-r}$. When composed with the translation automorphisms τ_a, $a\in\mathbb{C}$ given by $\tau_a T(u) = T(u-a)$, that is

$$\tau_a T^{(r)} = \delta_{r0} + \sum_{s=1}^{r} T^{(s)} \binom{r-1}{r-s} a^{r-s}$$

this yields a one–parameter family of evaluation homomorphisms $\mathrm{ev}_a = \mathrm{ev}\circ\tau_a$ given by

$$\mathrm{ev}_a(t_{ij}^{(r)}) = \delta_{r0}\delta_{ij} + E_{ij}a^{r-1} \tag{5.10}$$

5.13. **Hopf algebra structure.** $Y(\mathfrak{gl}_n)$ is a Hopf algebra with coproduct

$$\Delta(t_{ij}(u)) = \sum_{k=1}^{n} t_{ik}(u)\otimes t_{kj}(u)$$

For any $m\geq 2$, let $\Delta^{(m)} : Y(\mathfrak{gl}_n)\rightarrow Y(\mathfrak{gl}_n)^{\otimes m}$ denote the corresponding iterated coproduct. Then,

$$\Delta^{(m)}(t_{ij}^{(2)}) = \sum_{p=1}^{m}(t_{ij}^{(2)})_p + \sum_{\substack{1\leq k\leq n\\ 1\leq p<q\leq m}} (E_{ik})_p(E_{kj})_q \tag{5.11}$$

where $X_p = 1^{\otimes(p-1)}\otimes X\otimes 1^{\otimes(m-p)}$.

5.14. **Evaluation modules.** For any $\underline{a} = (a_1,\ldots,a_m)\in\mathbb{C}^m$ define $\mathrm{ev}_{\underline{a}} : Y(\mathfrak{gl}_n)\rightarrow U\mathfrak{gl}_n^{\otimes m}$ by

$$\mathrm{ev}_{\underline{a}} = \mathrm{ev}_{a_1}\otimes\cdots\otimes\mathrm{ev}_{a_m}\circ\Delta^{(m)}$$

Proposition. *The image of the trigonometric Casimir connection of* $\mathfrak{gl}_n$ *under the homomorphism* $\mathrm{ev}_{\underline{a}}$ *is the* $U\mathfrak{gl}_n^{\otimes m}$*–valued connection given by*

$$\nabla_{\underline{a}} = d - \sum_{i<j}\frac{d(\theta_i-\theta_j)}{e^{\theta_i-\theta_j}-1}\Delta^{(m)}(\kappa_{\theta_i-\theta_j}) - \sum_{i=1}^{n} d\theta_i\, D_{i,\underline{a}}$$

where

$$D_{i,\underline{a}} = 2\sum_{p=1}^{m} a_p(E_{ii})_p + 2\sum_{\substack{1\leq j\leq n\\ 1\leq p<q\leq m}}(E_{ij})_p(E_{ji})_q - \sum_{j<i}\Delta^{(m)}(\kappa_{\theta_j-\theta_i}) - \Delta^{(m)}(E_{ii}^2)$$

Proof. By construction $D_{i,\underline{a}} = \mathrm{ev}_{\underline{a}}(D_i)$ and is given by the above expression by (5.5), (5.11) and (5.10). ■

5.15. **The trigonometric dynamical differential equations for $\mathfrak{gl}_n$.** In [TV], Tarasov and Varchenko considered differential operators $\mathcal{D}_1, \ldots, \mathcal{D}_n$ in the variables $z_1, \ldots, z_n \in \mathbb{C}^\times$ with coefficients in $U\mathfrak{gl}_n^{\otimes m}$ given by

$$\mathcal{D}_i = z_i \partial_{z_i} + \lambda L_i(\underline{a}, \underline{z})$$

where $\lambda \in \mathbb{C}$, $\underline{a} = (a_1, \ldots, a_m) \in \mathbb{C}^m$, $\underline{z} = (z_1, \ldots, z_n)$ and

$$\begin{aligned} L_i(\underline{a}, \underline{z}) = \frac{\Delta^{(m)}(E_{ii}^2)}{2} &- \sum_{p=1}^{m} a_p (E_{ii})_p \\ &- \sum_{\substack{1 \leq j \leq n \\ 1 \leq p < q \leq m}} (E_{ij})_p (E_{jk})_q - \sum_{j \neq i} \frac{z_j}{z_i - z_j} \Delta^{(m)}(E_{ij}E_{ji} - E_{ii}) \end{aligned}$$

Set $z_i = e^{\theta_i}$ so that $z_i \partial_{z_i} = \partial_{\theta_i}$ and

$$\frac{z_j}{z_i - z_j} = \frac{1}{e^{\theta_i - \theta_j} - 1} = -\left(\frac{1}{e^{\theta_j - \theta_i} - 1} + 1 \right)$$

Since

$$E_{ij}E_{ji} - E_{ii} = \frac{1}{2}(\kappa_{\theta_i - \theta_j} - (E_{ii} + E_{jj}))$$

the operators $\mathcal{D}_i$ are the covariant derivatives for the connection

$$\nabla'_{\underline{a}} = d - \frac{\lambda}{2} \left(\sum_{i<j} \frac{d(\theta_i - \theta_j)}{e^{\theta_i - \theta_j} - 1} \Delta^{(m)}(\kappa_{\theta_i - \theta_j} - (E_{ii} + E_{jj})) + \sum_{i=1}^{n} d\theta_i \, D'_{i,\underline{a}} \right)$$

where

$$\begin{aligned} D'_{i,\underline{a}} = -\Delta^{(m)}(E_{ii}^2) + 2 \sum_{p=1}^{m} a_p (E_{ii})_p + 2 &\sum_{\substack{1 \leq j \leq n \\ 1 \leq p < q \leq m}} (E_{ij})_p (E_{ji})_q \\ &- \sum_{j<i} \Delta^{(m)}(\kappa_{\theta_j - \theta_i} - (E_{ii} + E_{jj})) \end{aligned}$$

By Proposition 5.14, $\nabla'_{\underline{a}}$ is the image of the trigonometric Casimir connection for $\mathfrak{gl}_n$[5] under the homomorphism $\mathrm{ev}_{\underline{a}} : Y(\mathfrak{gl}_n) \to (U\mathfrak{gl}_n)^{\otimes m}$ plus the $\mathfrak{h}$–valued, closed one–form

$$\frac{\lambda}{2} \Delta^{(m)} \left(\sum_{i<j} \frac{d(\theta_i - \theta_j)}{e^{\theta_i - \theta_j} - 1} (E_{ii} + E_{jj}) - \sum_{i} d\theta_i \sum_{j<i} (E_{ii} + E_{jj}) \right)$$

---

[5] when the latter is scaled by a factor of $\lambda/2$, as in Remark 3.8.

6. Bispectrality

We show in this section that the trigonometric Casimir connection with values in a tensor product of $Y(\mathfrak{g})$–modules commutes with the qKZ difference equations of Frenkel–Reshetikhin determined by the rational R–matrix of $Y(\mathfrak{g})$. This was checked by Tarasov–Varchenko for $\mathfrak{g} = \mathfrak{gl}_n$ in the case where all representations are evaluation modules [TV].

6.1. **Hopf algebra structure** [Dr1]**.** If $\mathfrak{g}$ is simple, $Y(\mathfrak{g})$ is a Hopf algebra with coproduct $\Delta : Y(\mathfrak{g}) \to Y(\mathfrak{g}) \otimes Y(\mathfrak{g})$ given on generators by

$$\Delta(x) = x \otimes 1 + 1 \otimes x$$

$$\Delta(J(x)) = J(x) \otimes 1 + 1 \otimes J(x) + \frac{\hbar}{2}[x \otimes 1, t]$$

where $t = \sum_a x_a \otimes x^a \in (\mathfrak{g}\otimes\mathfrak{g})^{\mathfrak{g}}$, with $\{x_a\}, \{x^a\}$ dual bases of $\mathfrak{g}$ with respect to the given inner product. Thus, if $\Delta^{(n)} : Y(\mathfrak{g}) \to Y(\mathfrak{g})^{\otimes n}$ is the iterated coproduct, then

$$\Delta^{(n)}(x) = \sum_{i=1}^{n} x^{(i)} \tag{6.1}$$

$$\Delta^{(n)}(J(x)) = \sum_{i=1}^{n} J(x)^{(i)} + \frac{\hbar}{2} \sum_{1\leq i<j\leq n} [x^{(i)}, t^{ij}] \tag{6.2}$$

where $x^{(i)} = 1^{\otimes(i-1)} \otimes x \otimes 1^{\otimes(n-i)}$ and $t^{ij} = \sum_a x_a^{(i)} (x^a)^{(j)}$.

6.2. **Translation automorphisms** [Dr1]**.** $Y(\mathfrak{g})$ possesses a one–parameter group of Hopf algebra automorphisms T_v, $v \in \mathbb{C}$ given by

$$T_v x = x \qquad \text{and} \qquad T_v J(x) = J(x) + vx$$

If $v_1, \ldots, v_n \in \mathbb{C}$, we set $T_{v_1,\ldots,v_n} = T_{v_1} \otimes \cdots \otimes T_{v_n} \in \mathrm{Aut}(Y(\mathfrak{g})^{\otimes n})$ and

$$\Delta_{v_1,\ldots,v_n} = T_{v_1,\ldots,v_n} \circ \Delta^{(n)} : Y(\mathfrak{g}) \to Y(\mathfrak{g})^{\otimes n} \tag{6.3}$$

6.3. **The universal R–matrix of** $Y(\mathfrak{g})$ [Dr1]**.** Let $R(u) \in Y(\mathfrak{g})\otimes Y(\mathfrak{g})[[u^{-1}]]$ be the universal R–matrix of $Y(\mathfrak{g})$. $R(u)$ satisfies

$$\Delta \otimes \mathrm{id}(R(u)) = R^{13}(u)R^{23}(u) \tag{6.4}$$

$$\mathrm{id} \otimes \Delta(R(u)) = R^{13}(u)R^{12}(u) \tag{6.5}$$

$$\Delta^{21}(x) = R(u)\Delta(x)R(u)^{-1} \tag{6.6}$$

$$T_{v,w}R(u) = R(u + v - w) \tag{6.7}$$

The above relations imply that R satisfies the quantum Yang–Baxter equations (QYBE) with spectral parameter

$$R^{12}(u)R^{13}(u + v)R^{23}(v) = R^{23}(v)R^{13}(u + v)R^{12}(u)$$

and the more general form of (6.6)

$$R(u)(T_{v,w}\Delta(x))R(u)^{-1} = T_{v,w}\Delta^{21}(x)$$

6.4. **The rational qKZ equations** [FR]**.** Let $V_1, \ldots, V_n$ be $Y(\mathfrak{g})$–modules and $d_i \in GL(V_i)$ be such that $d_i d_j R^{ij}(u) = R^{ij}(u) d_i d_j$ for any $1 \leq i < j \leq n$. Fix a step $\varkappa \in \mathbb{C}^\times$, let $a_1, \ldots, a_n \in \mathbb{C}$ be distinct and define operators

$$A_i = A_i(a_1, \ldots, a_n) \in \mathrm{End}(V_1 \otimes \cdots \otimes V_n)$$

for $i = 1, \ldots, n$ by

$$\begin{aligned} A_i = {} & R^{i-1\,i}(a_{i-1} - a_i - \varkappa)^{-1} R^{i-2\,i}(a_{i-2} - a_i - \varkappa)^{-1} \cdots R^{1\,i}(a_1 - a_i - \varkappa)^{-1} \cdot \\ & d_i \cdot R^{i\,n}(a_i - a_n) R^{i\,n-1}(a_i - a_{n-1}) \cdots R^{i\,i+1}(a_i - a_{i+1}) \end{aligned}$$

The (rational) qKZ equations of Frenkel–Reshetikhin are the system of difference equations $T_i f = A_i f$ where f takes values in $V_1 \otimes \cdots \otimes V_n$ and

$$T_i f(a_1, \ldots, a_n) = f(a_1, \ldots, a_{i-1}, a_i + \varkappa, a_{i+1}, \ldots, a_n)$$

They are a consistent system in that $[\not{T}_i, \not{T}_j] = 0$, where $\not{T}_i = A_i^{-1} T_i$ are the covariant difference operators.

6.5.

Lemma. *The following holds for any* $i = 1, \ldots, n$

$$\not{T}_1 \not{T}_2 \cdots \not{T}_i = (\widetilde{A}_i)^{-1} T_1 \cdots T_i$$

where $\widetilde{A}_i = \widetilde{A}_i(a_1, \ldots, a_n)$ *is given by*

$$\begin{aligned} \widetilde{A}_i = d_1 \cdots d_i & \big(R^{1\,n}(a_1 - a_n) \cdots R^{i\,n}(a_i - a_n)\big) \\ & \big(R^{1\,n-1}(a_1 - a_{n-1}) \cdots R^{i\,n-1}(a_i - a_{n-1})\big) \cdots \\ & \cdots \big(R^{1\,i+1}(a_1 - a_{i+1}) \cdots R^{i\,i+1}(a_i - a_{i+1})\big) \end{aligned} \tag{6.8}$$

Thus, $\widetilde{A}_n = d_1 \cdots d_n$ *and for any* $i \leq n-1$

$$\widetilde{A}_i = d_1 \cdots d_i\, \Delta^{(i)}_{a_1 - a_i, \ldots, a_{i-1} - a_i, 0} \otimes \Delta^{(n-i)}_{a_{i+1} - a_n, \ldots, a_{n-1} - a_n, 0} (R(a_i - a_n)) \tag{6.9}$$

Proof. Clearly,

$$\not{T}_1 \not{T}_2 \cdots \not{T}_i = \left(T_1 \cdots T_{i-1}(A_i) \cdots T_1(A_2) A_1\right)^{-1} T_1 \cdots T_i$$

and, for any $j \leq i$,

$$T_1 \cdots T_{j-1}(A_j) = (R^{i-1\,i})^{-1} (R^{i-2\,i})^{-1} \cdots (R^{1\,i})^{-1} \cdot d_i \cdot R^{i\,n} R^{i\,n-1} \cdots R^{i\,i+1}$$

where R^{kl} is shorthand for $R^{kl}(a_k - a_l)$. The first claimed identity now follows by induction on i using the QYBE. The second follows from the relations (6.4)–(6.5) and (6.7). ∎

6.6. **Bispectrality.** To couple the qKZ and trigonometric Casimir connection equations with values in the tensor product $V_1 \otimes \cdots \otimes V_n$, assume that each V_i is an integrable $\mathfrak{g}$–module and that d_i is the $GL(V_i)$–valued function on the torus H given by

$$d_i(e^u) = (e^{-u})^{(i)} \tag{6.10}$$

Let also $\nabla_{\underline{a}}$ be the trigonometric Casimir connection with values in the $Y(\mathfrak{g})$–module $T^*_{-a_1,\ldots,-a_n} V_1 \otimes \cdots \otimes V_n$ and scaled by a factor of $2\varkappa$ as in Remark 3.8. Thus, $\nabla_{\underline{a}}$ is the $\operatorname{End}(V_1 \otimes \cdots \otimes V_n)$–valued connection given by

$$\nabla_{\underline{a}} = d - \frac{1}{2\varkappa}\, \Delta^{(n)}_{a_1,\ldots,a_n}(B)$$

where

$$B = \frac{\hbar}{2} \sum_{\alpha \in \Phi_+} \frac{e^\alpha + 1}{e^\alpha - 1} d\alpha\, \kappa_\alpha - 2du_i\, J(u^i)$$

Theorem. *The qKZ operators $\not{T}_i$ commute with the trigonometric Casimir connection $\nabla_{\underline{a}}$.*

Proof. It suffices to prove that $[\nabla_{\underline{a}}, \not{T}_1 \cdots \not{T}_i] = 0$ for any $i = 1, \ldots, n$. Since

$$\begin{aligned}
&[d - (2\varkappa)^{-1}\Delta^{(n)}_{a_1,\ldots,a_n}(B), (\widetilde{A}_i)^{-1} T_1 \cdots T_i]\, (T_1 \cdots T_i)^{-1} \\
&\qquad = d\widetilde{A}_i^{-1} - (2\varkappa)^{-1} \widetilde{A}_i^{-1} (\mathrm{id} - \operatorname{Ad}(T_1 \cdots T_i)) \Delta^{(n)}_{a_1,\ldots,a_n}(B) \\
&\qquad\qquad - (2\varkappa)^{-1} [\Delta^{(n)}_{a_1,\ldots,a_n}(B), (\widetilde{A}_i)^{-1}]
\end{aligned}$$

the claim follows from the two lemmas below. ■

6.7.

Lemma.

$$(d\widetilde{A}_i)\widetilde{A}_i^{-1} = (2\varkappa)^{-1}(\operatorname{Ad}(T_1 \cdots T_i) - \mathrm{id})\Delta^{(n)}_{a_1,\ldots,a_n}(B) \tag{6.11}$$

Proof. By (6.8), the left–hand side of (6.11) is equal to

$$d(d_1 \cdots d_i)\,(d_1 \cdots d_i)^{-1} = -\sum_{j=1}^{i} du_a (u^a)^{(j)}$$

where we used (6.10). Write $B = B_1 + B_2$ where

$$B_1 = \frac{\hbar}{2} \sum_{\alpha \in \Phi_+} d\alpha\, \frac{e^\alpha + 1}{e^\alpha - 1} \kappa_\alpha \qquad \text{and} \qquad B_2 = -2du_i\, J(u^i)$$

Since B_1 takes values in $U\mathfrak{g}$, $\Delta_{a_1,\ldots,a_n}(B_1)^{(n)}$ is independent of $a_1, \ldots, a_n$ and the right–hand side of (6.11) is equal to $(2\varkappa)^{-1}(\operatorname{Ad}(T_1 \cdots T_i) - \mathrm{id})\Delta^{(n)}_{a_1,\ldots,a_n}(B_2)$. By (6.2), for any $x \in \mathfrak{g}$,

$$(\operatorname{Ad}(T_1 \cdots T_i) - \mathrm{id})\Delta^{(n)}_{a_1,\ldots,a_n}(J(x)) = \sum_{j=1}^{i} \varkappa x^{(i)}$$

so that the right–hand side of (6.11) is equal to $-\sum_{j=1}^{i} du_a (u^a)^{(j)}$. ■

6.8.

Lemma.

$$[\widetilde{A}_i, \Delta^{(n)}_{a_1,\ldots,a_n}(B)] = 0$$

Proof. For any $x \in Y(\mathfrak{g})$ and $1 \leq i \leq n$,

$$\Delta^{(n)}_{a_1,\ldots,a_n}(x) = \Delta^{(i)}_{a_1-a_i,\ldots,a_{i-1}-a_i,0} \otimes \Delta^{(n-i)}_{a_{i+1}-a_n,\ldots,a_{n-1}-a_n,0} \circ \Delta_{a_i,a_n}(x)$$

so that, by (6.9) and the fact that $d_1 \cdots d_i = \Delta^{(i)}(d_1)$, it suffices to prove the claimed identity for $n = 2$ and $i = 1$. We have

$$\begin{aligned} & d_1^{-1} \, [d_1 R(a_1 - a_2), \Delta_{a_1,a_2}(B)] \, R(a_1 - a_2)^{-1} \\ & \quad = (\mathrm{id} - \mathrm{Ad}(d_1^{-1}))\Delta_{a_1,a_2}(B) + (\mathrm{Ad}(R(a_1 - a_2)) - \mathrm{id})\Delta_{a_1,a_2}(B) \end{aligned} \tag{6.12}$$

Let $u \in \mathfrak{h}$. By (6.2),

$$(\mathrm{id} - \mathrm{Ad}(d_1^{-1}))\Delta_{a_1,a_2}(J(u)) = \frac{\hbar}{2}(\mathrm{id} - \mathrm{Ad}(d_1^{-1}))[u^{(1)}, t]$$

and, by (6.6),

$$(\mathrm{Ad}(R(a_1 - a_2)) - \mathrm{id})\Delta_{a_1,a_2}(J(u)) = \frac{\hbar}{2}\,[u^{(2)} - u^{(1)}, t] = -\hbar\,[u^{(1)}, t]$$

Thus, the right–hand side of (6.12) with B replaced by $B_2 = -2du_a\, J(u^a)$ is equal to

$$\hbar\, du_a (\mathrm{id} + \mathrm{Ad}(d_1^{-1}))[u^{a(1)}, t]$$

Since $R(a_1 - a_2)$ commutes with $\Delta_{a_1,a_2}(B_1)$, we have left to compute

$$(\mathrm{id} - \mathrm{Ad}(d_1^{-1}))\Delta_{a_1,a_2}(B_1) = \hbar \sum_{\alpha \in \Phi_+} d\alpha\, \frac{e^\alpha + 1}{e^\alpha - 1}(\mathrm{id} - \mathrm{Ad}(d_1^{-1}))\overline{t}_\alpha$$

where $\overline{t}_\alpha = x_\alpha \otimes x_{-\alpha} + x_{-\alpha} \otimes x_\alpha$, with $x_{\pm\alpha} \in \mathfrak{g}_{\pm\alpha}$ such that $(x_\alpha, x_{-\alpha}) = 1$, so that $\Delta(\kappa_\alpha) = \kappa_\alpha \otimes 1 + 1 \otimes \kappa_\alpha + 2\overline{t}_\alpha$. By (6.10),

$$(\mathrm{id} - \mathrm{Ad}(d_1^{-1}))\overline{t}_\alpha = (1 - e^\alpha)(x_\alpha \otimes x_{-\alpha} - e^{-\alpha} x_{-\alpha} \otimes x_\alpha)$$

so that, for any $\alpha \in \Phi_+$ and $u \in \mathfrak{h}$,

$$\begin{aligned} & \alpha(u)\frac{e^\alpha + 1}{e^\alpha - 1}(\mathrm{id} - \mathrm{Ad}(d_1^{-1}))\overline{t}_\alpha \\ & \qquad = -\alpha(u)\left((e^\alpha + 1)x_\alpha \otimes x_{-\alpha} - (e^{-\alpha} + 1)x_{-\alpha} \otimes x_\alpha\right) \\ & \qquad\qquad = -[u^{(1)}, (\mathrm{id} + \mathrm{Ad}(d_1^{-1}))\overline{t}_\alpha] \end{aligned}$$

whence the claimed result. ■

Remark. The proof of Theorem 6.6 works almost *verbatim* for $\mathfrak{g} = \mathfrak{gl}_n$ and gives the commutation of the rational qKZ connection and trigonometric Casimir connections for $Y(\mathfrak{gl}_n)$.

7. The affine KZ connection

We show in this section that the degenerate affine Hecke algebra $\mathcal{H}'$ of W is, very roughly speaking, the 'Weyl group' of the Yangian $Y(\mathfrak{g})$. More precisely, we show that if V is a $Y(\mathfrak{g})$–module whose restriction to $\mathfrak{g}$ is *small*, the canonical action of W on the zero weight space $V[0]$ extends to one of $\mathcal{H}'$. Moreover, the trigonometric Casimir connection with values in $V[0]$ coincides with Cherednik's affine KZ connection with values in this $\mathcal{H}'$–module.

7.1. **The degenerate affine Hecke algebra.** Let K be the vector space of W–invariant functions $\Phi \to \mathbb{C}$ and denote the natural linear coordinates on K by k_α, $\alpha \in \Phi/W$. Recall [Lu] that the degenerate affine Hecke algebra $\mathcal{H}'$ associated to W is the algebra over $\mathbb{C}[K]$ generated by the group algebra $\mathbb{C}W$ and the symmetric algebra $S\mathfrak{h}$ subject to the relations

$$s_i x_u - x_{s_i(u)} s_i = k_{\alpha_i} \alpha_i(u) \tag{7.1}$$

for any simple reflection $s_i \in W$ and linear generator x_u, $u \in \mathfrak{h}$, of $S\mathfrak{h}$.

7.2. **The affine KZ connection.** The AKZ connection is the trigonometric, $\mathcal{H}'$–valued connection given by

$$\nabla = d - \sum_{\alpha \in \Phi_+} \frac{d\alpha}{e^\alpha - 1} k_\alpha s_\alpha - du_i \, x_{u^i} \tag{7.2}$$

where $\{u_i\}$, $\{u^i\}$ are dual bases of $\mathfrak{h}^*$, $\mathfrak{h}$ respectively. This connection was defined by Cherednik in [Ch1, Ch2] and proved to be flat and W–equivariant. This may also be obtained as a consequence of Theorem 2.23. Indeed, the relations (tt), with $t_\alpha = k_\alpha s_\alpha$ are easily verified and, as pointed out by Cherednik, the remaining relations are precisely those defining $\mathcal{H}'$.

Remark. The δ–form (2.7) of the AKZ connection corresponds to Drinfeld's presentation of $\mathcal{H}'$ in terms of $\mathbb{C}W$ and non–commuting elements y_u which transform like the reflection representation of W (see [Dr3] and [RS]).[6]Indeed, it is given by

$$\nabla = d - \frac{1}{2} \sum_{\alpha \in \Phi_+} \frac{e^\alpha - 1}{e^\alpha - 1} d\alpha \, k_\alpha s_\alpha - du_i \, y_{u^i}$$

where the elements y_u are defined by (2.6) as

$$y_u = x_u - \frac{1}{2} \sum_{\alpha \in \Phi_+} \alpha(u) k_\alpha s_\alpha \tag{7.3}$$

and therefore satisfy $s_i \, y_u s_i = y_{s_i(u)}$ by Proposition 2.22.

---

[6] I owe this observation to Pavel Etingof.

7.3. W**–action on zero weight spaces of** $\mathfrak{g}$**–modules.** Let G be the complex, simply–connected Lie group with Lie algebra $\mathfrak{g}$, H the maximal torus with Lie algebra $\mathfrak{h}$ and $N(H) \subset G$ its normaliser. If V is an integrable $\mathfrak{g}$–module, the action of $N(H)$ on V permutes the weight spaces compatibly with the action of $N(H)$ on H. In particular, it acts on the zero weight space $V[0]$ and this action factors through $W = N(H)/H$.

7.4. **Small** $\mathfrak{g}$**–modules.** Recall that a $\mathfrak{g}$–module V is *small* if 2α is not a weight of V for any root α [Br, Re, Re2]. If V is a small $\mathfrak{g}$–module with a non–trivial zero weight space $V[0]$, the restriction to $V[0]$ of the square e_α^2 of a raising operator maps to the weight space $V[2\alpha]$ and is therefore zero. This implies the following result [TL2, Prop. 9.1]

Lemma. *If V is an integrable, small $\mathfrak{g}$–module, the following holds on the zero weight space $V[0]$*

$$\kappa_\alpha = (\alpha, \alpha)(1 - s_\alpha)$$

where the right–hand side refers to the action of the reflection $s_\alpha \in W$ on $V[0]$.

7.5. Let $\mathcal{H}'_\hbar$ be the degenerate affine Hecke algebra of W with parameters

$$k_\alpha = -\hbar(\alpha, \alpha) \tag{7.4}$$

Theorem. *Let V be a finite–dimensional $Y(\mathfrak{g})$–module whose restriction to $\mathfrak{g}$ is small.*

1. *The canonical W–action on the zero weight space $V[0]$ together with either of the equivalent assignments*

$$x_u \to -2T(u)_1 + \frac{1}{2}\sum_{\alpha\in\Phi_+} k_\alpha \alpha(u) \tag{7.5}$$

$$y_u \to -2J(u) \tag{7.6}$$

   yield an action of $\mathcal{H}'_\hbar$ on $V[0]$.

2. *The trigonometric Casimir connection of $\mathfrak{g}$ with values in* $\mathrm{End}(V[0])$ *is equal to the sum of the AKZ connection with values in the $\mathcal{H}'_\hbar$–module $V[0]$ and the scalar valued one–form*

$$\mathcal{A} = \frac{1}{2}\sum_{\alpha\in\Phi} k_\alpha \frac{d\alpha}{e^\alpha - 1} \tag{7.7}$$

Proof. The trigonometric Casimir connection with values in $\mathrm{End}(V[0])$ reads, by Lemma 7.4

$$\begin{aligned}\nabla &= d - \hbar \sum_{\alpha\in\Phi_+} \frac{d\alpha}{e^\alpha - 1}(\alpha, \alpha)(1 - s_\alpha) + 2du_i\, T(u^i)_1 \\ &= d - \sum_{\alpha\in\Phi_+} \frac{d\alpha}{e^\alpha - 1} k_\alpha s_\alpha + 2du_i\, T(u^i)_1 + \sum_{\alpha\in\Phi_+} k_\alpha \frac{d\alpha}{e^\alpha - 1}\end{aligned}$$

where the weights k_α are given by (7.4). By (2.4)

$$\sum_{\alpha\in\Phi}(\alpha,\alpha)\frac{d\alpha}{e^\alpha-1}=2\sum_{\alpha\in\Phi_+}(\alpha,\alpha)\frac{d\alpha}{e^\alpha-1}+\sum_{\alpha\in\Phi_+}(\alpha,\alpha)d\alpha$$

so that if $\mathcal{A}$ is given by (7.7), then

$$\nabla-\mathcal{A}=d-\sum_{\alpha\in\Phi_+}k_\alpha\frac{d\alpha}{e^\alpha-1}s_\alpha+du_i\left(2T(u^i)_1-\frac{1}{2}\sum_{\alpha\in\Phi_+}k_\alpha\alpha(u^i)\right)$$

Applying Proposition 2.22 to $\nabla-\mathcal{A}$ which is W–equivariant since ∇ and $\mathcal{A}$ are, shows that the map (7.5) gives an action of $\mathcal{H}'_\hbar$ on $V[0]$. This proves (1) and (2). The equivalence of (7.5) and (7.6) follows easily from §3.3 and (7.3). ■

Remark. Theorem 7.5 extends to the trigonometric setting the relation between the rational Casimir and KZ connections proved in [TL2, Prop. 9.1].

7.6. **The adjoint representation.** Drinfeld proved that, for any simple $\mathfrak{g}$, the direct sum $\overline{\mathfrak{g}}=\mathfrak{g}\oplus\mathbb{C}$ of the adjoint and trivial representations of $\mathfrak{g}$ admits an extension to an action of $Y(\mathfrak{g})$ on $\overline{\mathfrak{g}}$ [Dr2, Thm. 8]. It is easy to check that the corresponding action of $\mathcal{H}'_\hbar$ on $\overline{\mathfrak{g}}[0]=\mathfrak{h}\oplus\mathbb{C}$ given by Theorem 7.5 coincides with its action on affine linear functions on $\mathfrak{h}^*$ given by rational Dunkl operators (see, *e.g.* [Ki]).

7.7. **The case of $\mathfrak{sl}_n$.** Let $\mathfrak{g}=\mathfrak{sl}_n$ and $V=\mathbb{C}^n$ its vector representation. A simple inspection shows that $V^{\otimes n}$ is a small [Re2]. The zero weight space $V^{\otimes n}[0]$ possesses two actions of the symmetric group: one arising from the Weyl group action of $\mathfrak{S}_n$, the other from the permutation of the tensor factors, under which it identifies with the group algebra $\mathbb{C}\mathfrak{S}_n$.

The $\mathfrak{sl}_n$–module $V^{\otimes n}$ may be endowed with an action of $Y(\mathfrak{g})$ depending on $a_1,\ldots,a_n\in\mathbb{C}$ obtained by composing the coproduct $\Delta^{(n)}:Y(\mathfrak{g})\to Y(\mathfrak{g})^{\otimes n}$ with the evaluation homomorphisms $\mathrm{ev}_{a_i}:Y(\mathfrak{g})\to U\mathfrak{g}$. It is easy to check that the action of $\mathcal{H}'_\hbar$ on $V^{\otimes n}[0]$ given by Theorem 7.5 coincides with that on the induced representation $\mathrm{ind}_{S\mathfrak{h}}^{\mathcal{H}'}\mathbb{C}_{a_1,\ldots,a_n}$.

8. Appendix: Tits extensions of affine Weyl groups

In this appendix, we review the definition of the Tits extension $\widetilde{W}$ of a Weyl group W. We then define the reduced Tits extension $\widetilde{W}^{\mathrm{red}}$ of W and show that, when W is an affine Weyl group, $\widetilde{W}^{\mathrm{red}}$ is isomorphic to the semi–direct product of the Tits extension of the finite Weyl group underlying W by the corresponding coroot lattice (Theorem 8.10).

8.1. **Weyl groups and braid groups.** Let $A = (a_{ij})_{i,j\in\mathbf{I}}$ be a generalised Cartan matrix and $(\mathfrak{h}, \Delta, \Delta^\vee)$ its unique realisation. Thus, $\mathfrak{h}$ is a complex vector space of dimension $2|\mathbf{I}| - \operatorname{rank}(A)$, $\Delta = \{\alpha_i\}_{i\in\mathbf{I}} \subset \mathfrak{h}^*$ and $\Delta^\vee = \{\alpha_i^\vee\}_{i\in\mathbf{I}} \subset \mathfrak{h}$ are linearly independent sets and

$$\langle \alpha_i^\vee, \alpha_j\rangle = a_{ij}$$

Recall that the Weyl group $W = W(A)$ attached to A is the subgroup of $GL(\mathfrak{h}^*)$ generated by the reflections [Ka, §3.7]

$$s_i(\lambda) = \lambda - \langle \lambda, \alpha_i^\vee\rangle \alpha_i$$

or, equivalently, the subgroup of $GL(\mathfrak{h})$ generated by the dual reflections

$$s_i^\vee(t) = t - \langle t, \alpha_i\rangle \alpha_i^\vee$$

By [Ka, Prop. 3.13], the defining relations of W are

$$s_i^2 = 1$$
$$(s_i s_j)^{m_{ij}} = 1$$

where for any $i \neq j$, m_{ij} is equal to $2, 3, 4, 6$ or ∞ according to whether $a_{ij}a_{ji}$ is equal to $0, 1, 2, 3$ or ≥ 4.

The *braid group* $B = B(A)$ attached to A is the group with generators S_i, $i \in \mathbf{I}$ and relations

$$\underbrace{S_i S_j \cdots}_{m_{ij}} = \underbrace{S_j S_i \cdots}_{m_{ij}}$$

for any $i \neq j$.

8.2. **Tits extensions of Weyl groups.**

Definition ([Ti])**.** The *Tits extension* of W is the group $\widetilde{W}$ with generators $\widetilde{s}_i$, $i \in \mathbf{I}$ and relations

$$\underbrace{\widetilde{s}_i \widetilde{s}_j \cdots}_{m_{ij}} = \underbrace{\widetilde{s}_j \widetilde{s}_i \cdots}_{m_{ij}} \tag{8.1}$$

$$\widetilde{s}_i^4 = 1 \tag{8.2}$$

$$\widetilde{s}_i^2 \widetilde{s}_j^2 = \widetilde{s}_j^2 \widetilde{s}_i^2 \tag{8.3}$$

$$\widetilde{s}_i \widetilde{s}_j^2 \widetilde{s}_i^{-1} = \widetilde{s}_j^2 (\widetilde{s}_i^2)^{-a_{ji}} \tag{8.4}$$

8.3. Let $\mathfrak{g} = \mathfrak{g}(A)$ be the Kac–Moody algebra corresponding to the Cartan matrix A with generators $t \in \mathfrak{h}$ and e_i, f_i, $i \in \mathbf{I}$. Recall that a representation of $\mathfrak{g}$ is integrable if $\mathfrak{h} \subset \mathfrak{g}$ acts semi–simply with finite–dimensional eigenspaces and e_i, f_i act locally nilpotently. The next two results explain the relevance and structure of the Tits extension $\widetilde{W}$.

Proposition. *Let V be an integrable representation of $\mathfrak{g}$. Then, the triple exponentials*

$$r_i = \exp(e_i)\exp(-f_i)\exp(e_i)$$

are well–defined elements of $GL(V)$ *and the assignment* $\widetilde{s}_i \to r_i$ *yields a representation of* $\widetilde{W}$ *on* V *mapping* $\widetilde{s}_i^2$ *to* $\exp(\pi\sqrt{-1}\alpha_i^\vee)$.

Proof. The r_i are clearly well defined and satisfy

$$r_i \cdot t \cdot r_i^{-1} = s_i^\vee(t)$$

for any $t \in \mathfrak{h}$ and $r_i^2 = \exp(\pi\sqrt{-1}\alpha_i^\vee)$ [Ka, §3.8], from which (8.2)–(8.4) readily follow. Let now $i \neq j$ be such that $m_{ij} < \infty$. Then, the Lie subalgebra $\mathfrak{g}_{ij}$ of $\mathfrak{g}$ generated by $e_i, f_i, \alpha_i^\vee$ and $e_j, f_j, \alpha_j^\vee$ is finite–dimensional and semi–simple and V integrates to a representation of the complex, connected and simply–connected Lie group G_{ij} with Lie algebra $\mathfrak{g}_{ij}$. By [Ti], r_i and r_j satisfy the braid relations (8.1) when regarded as elements of G_{ij}[7], and these therefore hold in $GL(V)$. ■

8.4. Let $Q^\vee \subset \mathfrak{h}$ be the lattice spanned by the coroots $\alpha_i^\vee$, $i \in \mathbf{I}$.

Proposition ([Ti])**.** $\widetilde{W}$ *is an extension of* W *by the abelian group* Z *generated by the elements* $\widetilde{s}_i^2$. Z *is isomorphic, as* W*–module to* $Q^\vee/2Q^\vee \cong \mathbb{Z}_2^{|\mathbf{I}|}$.

Proof. Let $K \supset Z$ the kernel of the canonical projection $\widetilde{W} \to W$. By (8.4), Z is a normal subgroup of $\widetilde{W}$ and $\widetilde{W}/Z$ is generated by the images $\overline{s}_i$ of $\widetilde{s}_i$ which, in addition to the braid relations (8.1), satisfy $\overline{s}_i^2 = 1$. Thus, $\widetilde{W}/Z$ is a quotient of W, $K = Z$ and $\widetilde{W}/Z \cong W$. Note next that, by (8.2)–(8.4), the assignment $\alpha_i^\vee \to \widetilde{s}_i^2$ extends to a W–equivariant surjection $Q^\vee/2Q^\vee \to Z$. To prove that this is an isomorphism it suffices to exhibit, for any $i \in \mathbf{I}$ a $\mathbb{Z}_2$–valued character χ_i of Z such that $\chi_i(\widetilde{s}_j^2) = (-1)^{\delta_{ij}}$. Let λ_i be the ith fundamental weight of $\mathfrak{g}$, so that $\langle \lambda_i, \alpha_j^\vee \rangle = \delta_{ij}$, V_i the irreducible $\mathfrak{g}$–module with highest weight λ_i and $v_i \in V_i$ a nonzero highest weight vector. V_i is integrable and since $r_j^2 = \exp(\sqrt{-1}\pi\alpha_j^\vee)$, we have $r_j^2 v_i = (-1)^{\delta_{ij}} v_i$. ■

8.5. **Reduced Tits extensions.** For any $v = \sum_i m_i \alpha_i^\vee \in Q^\vee$, set

$$\widetilde{s}_v^2 = \prod_i (\widetilde{s}_i^2)^{m_i} \in Z$$

so that for any $w \in W$ and lift $\widetilde{w} \in \widetilde{W}$, $\widetilde{w}\widetilde{s}_v^2\widetilde{w}^{-1} = \widetilde{s}_{wv}^2$. By [Ka, Prop. 1.6], the center $\mathfrak{c}$ of $\mathfrak{g}$ is equal to

$$\mathfrak{c} = \{t \in \mathfrak{h} | \langle t, \alpha_i \rangle = 0 \text{ for any } i \in \mathbf{I}\} \tag{8.5}$$

The Weyl group operates trivially on $\mathfrak{c} \subset \mathfrak{h}$ and it follows from (8.4) that the subgroup $Z_{\mathfrak{c}} \subset Z$ generated by the elements $\widetilde{s}_v^2$, with $v \in \mathfrak{c} \cap Q^\vee$ lies in the centre of $\widetilde{W}$.

Definition. The *reduced Tits extension* $\widetilde{W}^{\text{red}}$ of W is the quotient

$$\widetilde{W}^{\text{red}} = \widetilde{W}/Z_{\mathfrak{c}}$$

---

[7] Tits' argument is reproduced in the proof of (i) of Proposition 8.9.

By Proposition 8.4, $\widetilde{W}^{\mathrm{red}}$ is an extension of W by $Q^\vee/(2Q^\vee + \mathfrak{c}\cap Q^\vee) \cong \mathbb{Z}_2^{\mathrm{rank}(A)}$.

8.6. **Reduced Tits extensions of affine Weyl groups.** Assume henceforth that $A = (a_{ij})_{0\leq i,j\leq n}$ is an affine Cartan matrix of untwisted type. Altering our notations, we denote by $\mathfrak{g}$ the underlying complex, semi–simple Lie algebra and by $\mathfrak{h}$, $\{\alpha_i\}_{i=1}^n$, $\{\alpha_i^\vee\}_{i=1}^n$, W and $Q^\vee$ its Cartan subalgebra, simple roots, simple coroots, Weyl group and coroot lattice respectively. Thus, for any $1 \leq i, j \leq n$,

$$a_{ij} = \langle \alpha_i^\vee, \alpha_j\rangle, \quad a_{0j} = -\langle \theta^\vee, \alpha_j\rangle \quad \text{and} \quad a_{j0} = -\langle \alpha_j^\vee, \theta\rangle$$

where $\theta \in \mathfrak{h}^*$ is the highest root of $\mathfrak{g}$. It is well known that the (affine) Weyl group W_a attached to A is isomorphic to the semi–direct product $W \ltimes Q^\vee$ [Ka, prop. 6.5]. The isomorphism is given by mapping s_i to $(s_i, 0)$ for $i \geq 1$ and s_0 to $(s_\theta, -\theta^\vee)$.

The subspace $\mathfrak{c}$ defined by (8.5) is spanned by the element

$$K = \alpha_0^\vee + \sum_{i=1}^n m_i\alpha_i^\vee$$

where the m_i are the positive integers such that $\theta^\vee = \sum_{i=1}^n m_i\alpha_i^\vee$ [Ka, Prop. 6.2]. It follows that the reduced Tits extension $\widetilde{W_a}^{\mathrm{red}}$ of W_a is the quotient of $\widetilde{W_a}$ by the relation

$$\widetilde{s}_0^2 \cdot \prod_{i=1}^n (\widetilde{s}_i^2)^{m_i} = 1 \tag{8.6}$$

8.7. **Loop groups.** The structure of the reduced Tits extension of W_a will be determined in §8.7–§8.10 by embedding $\widetilde{W_a}^{\mathrm{red}}$ into the loop group corresponding to $\mathfrak{g}$.

Let $L\mathfrak{g} = \mathfrak{g}[z, z^{-1}]$ be the loop algebra of $\mathfrak{g}$ and d the derivation of $L\mathfrak{g}$ defined by $dx(m) = mx(m)$, where $x(m) = x \otimes z^m$. Then, $L\mathfrak{g} \rtimes \mathbb{C}d$ is the quotient of the Kac–Moody algebra corresponding to A by the central element K defined above. Let G be the complex, connected and simply connected Lie group with Lie algebra $\mathfrak{g}$ and $LG = G(\mathbb{C}[z, z^{-1}])$ the group of polynomial loops into G. Let $H \subset G$ be the maximal torus with Lie algebra $\mathfrak{h}$. The group $\mathbb{C}^*$ acts on LG by reparametrisation fixing $G \supset H$ and $H \times \mathbb{C}^*$ is a maximal abelian subgroup of the semi–direct product $LG \rtimes \mathbb{C}^*$. By [PS], Prop. 5.2, the normaliser of $H \times \mathbb{C}^*$ in $LG \rtimes \mathbb{C}^*$ is equal to $(N(H) \ltimes H^\vee) \rtimes \mathbb{C}^*$ where $N(H)$ is the normaliser of H in G and $H^\vee = \mathrm{Hom}_{\mathbb{Z}}(\mathbb{C}^*, H) \subset LG$ is isomorphic to the coroot lattice $Q^\vee$ by

$$\lambda \in Q^\vee \longrightarrow \Big(z \to z^\lambda = \exp_H(-\ln(z)\lambda)\Big)$$

The quotient $N(H \times \mathbb{C}^*)/H \times \mathbb{C}^*$ is therefore isomorphic to the affine Weyl group $W_a = W \ltimes H^\vee$.

8.8. For each real root $\widetilde{\alpha} = (\alpha, n)$ of LG, the subalgebra $\mathfrak{sl}_2^{\widetilde{\alpha}}$ of $L\mathfrak{g}$ spanned by

$$e_{\widetilde{\alpha}} = e_\alpha(n), \quad f_{\widetilde{\alpha}} = f_\alpha(-n) \quad \text{and} \quad h_\alpha$$

is the Lie algebra of a closed subgroup of LG isomorphic to $SL_2(\mathbb{C})$. This is obvious if $n = 0$ and follows in the general case from the fact that $\mathfrak{sl}_2^{(\alpha,n)}$ is conjugate to $\mathfrak{sl}_2^{(\alpha,0)}$. Indeed, any element γ_λ of the coweight lattice $\mathrm{Hom}(\mathbb{C}^*, H/Z) \subset L(G/Z)$ induces by conjugation an automorphism of LG such that

$$\mathrm{Ad}(\gamma_\lambda) e_\alpha(n) = e_\alpha(n - \langle \lambda, \alpha \rangle) \qquad \text{and} \qquad \mathrm{Ad}(\gamma_\lambda) f_\alpha(n) = f_\alpha(n + \langle \lambda, \alpha \rangle)$$

8.9. Let now $\alpha_i = (\alpha_i, 0)$, $i = 1 \ldots n$ and $\alpha_0 = (-\theta, 1)$ be the simple roots of LG. For each $i = 0 \ldots n$, let $SL_2(\mathbb{C}) \cong G_i \subset LG$ be the corresponding subgroup, $H_i \subset G_i$ its torus and N_i the normaliser of H_i in G_i. Note that any element of $N_i \setminus H_i$ is of the form

$$\exp(e_i) \exp(-f_i) \exp(e_i) = \exp(-f_i) \exp(e_i) \exp(-f_i)$$

for some choice of root vectors $e_i \in (L\mathfrak{g})_{\alpha_i}, f_i \in (L\mathfrak{g})_{-\alpha_i}$ such that $[e_i, f_i] = \alpha_i^\vee$ if $i \geq 1$ and $-\theta^\vee$ if $i = 0$.

Let B_a be the (affine) braid group corresponding to A and $S_0, S_1, \ldots, S_n$ its generators.

Proposition.

(1) *For any choice of $\sigma_i \in N_i \setminus H_i$, $i = 0 \ldots n$, the assignment $S_i \to \sigma_i$ extends uniquely to a homomorphism $\sigma : B_a \to N(H) \ltimes H^\vee$.*

(2) *σ factors through an isomorphism of the reduced Tits extension $\widetilde{W_a}^{\mathrm{red}}$ onto its image in $N(H) \ltimes H^\vee$.*

(3) *If $\sigma, \sigma' : \widetilde{W_a}^{\mathrm{red}} \to N(H) \ltimes H^\vee$ are the homomorphisms corresponding to the choices $\{\sigma_i\}$ and $\{\sigma'_i\}$ respectively, there exists $t \in H \times \mathbb{C}^*$ such that, for any $\widetilde{s} \in \widetilde{W_a}^{\mathrm{red}}$, $\sigma(\widetilde{s}) = t \sigma'(\widetilde{s}) t^{-1}$.*

Proof. (1) the following argument is due to Tits [Ti]. Let $i \neq j$ be such that m_{ij} is finite and set $s_{ij} = s_i s_j \cdots \in W_a$ and $\sigma_{ij} = \sigma_i \sigma_j \cdots \in N(H) \ltimes H^\vee$ where each product has $m_{ij} - 1$ factors. The braid relations in W_a may be written as $s_{ij} s_{j'} = s_j s_{ij}$ where $j' = j$ or i according to whether m_{ij} is even or odd. Thus, $s_{ij}^{-1} s_j s_{ij} = s_{j'}$ and therefore,

$$\Delta_{ij} = \sigma_{j'}^{-1} \sigma_{ij}^{-1} \sigma_j \sigma_{ij} \in H \cap \left(\sigma_{j'}^{-1} \sigma_{ij}^{-1} N_j \sigma_{ij} \right) = H \cap \sigma_{j'}^{-1} N_{j'} = H_{j'}$$

Repeating the argument with i and j permuted, we find that $\Delta_{ji} \in H_{i'}$ with $i' = i$ or j according to whether m_{ij} is even or odd. Thus, $\Delta_{ij} = \Delta_{ji}^{-1} \in H_{i'} \cap H_{j'} = \{1\}$ where the latter assertion follows by follows from the simple connectedness of G.

(2) The σ_i satisfy (8.2)–(8.4) and (8.6) since, for any $x_j \in N_j \setminus H_j$, $x_j^2 = \exp(i\pi\alpha_j^\vee)$ for $j \geq 1$ and $x_0^2 = \exp(-i\pi\theta^\vee)$. Thus σ descends to $\widetilde{W_a}^{\mathrm{red}}$.

Since the diagram

$$\begin{tikzcd}
\widetilde{W_a}^{\mathrm{red}} \arrow[r] \arrow[rd] & N(H) \ltimes H^\vee \arrow[d] \\
 & W_a
\end{tikzcd}$$

is commutative, the kernel of σ is contained in $Z/Z_{\mathfrak{c}} \cong \mathbb{Z}_2^n$ and is therefore trivial since, due to the simple–connectedness of G, the subgroup of G generated by $\sigma_j^2 = \exp(\pi i \alpha_j^\vee)$, $j = 1 \ldots n$ is isomorphic to $\mathbb{Z}_2^n$.

(3) For $i = 1, \ldots, n$, let $t_i \in H_i$ be such that $\sigma_i = t_i \sigma_i'$ and choose $c_i \in \mathbb{C}$ such that $t_i = \exp(c_i h_{\alpha_i})$. Since $s_i \lambda_j^\vee = \lambda_j^\vee - \delta_{ij}\alpha_i^\vee$, where the $\lambda_j^\vee \in \mathfrak{h}$ are the fundamental coweights of $\mathfrak{g}$, we find, with $\overline{t} = \exp(\sum_{j=1}^n c_j \lambda_j^\vee) \in H$,

$$\overline{t} \cdot \sigma_i' \cdot \overline{t}^{-1} \exp(c_i h_{\alpha_i}) \cdot \sigma_i' = \sigma_i$$

Let now $t_0 = \exp(c_0 h_\theta) \in H_0$ be such that $\sigma_0 = t_0 \sigma_0'$. Since for any $x \in \mathbb{C}$,

$$\exp(xd)\sigma_0' \exp(-xd) = \exp(-x h_\theta)\sigma_0'$$

we find, with $y = \sum_j c_j \langle \lambda_j^\vee, \theta \rangle - c_0$, that

$$\overline{t} \exp(yd)\sigma_0' \exp(-yd)\overline{t}^{-1} = \exp((-y + \sum_j c_j \langle \lambda_j^\vee, \theta \rangle) h_\theta)\sigma_0' = \sigma_0$$

so that $t = \overline{t}\exp(yd)$ is the required element. ■

8.10. The following is the main result of this appendix.

Theorem. *The inclusion* $\widetilde{W} \hookrightarrow \widetilde{W_a}^{\mathrm{red}}$ *extends to an isomorphism* $\widetilde{W} \ltimes Q^\vee \to \widetilde{W_a}^{\mathrm{red}}$ *making the following a commutative diagram*

$$\begin{tikzcd}
\widetilde{W} \ltimes Q^\vee \arrow[r] \arrow[d] & \widetilde{W_a}^{\mathrm{red}} \arrow[d] \\
W \ltimes Q^\vee \arrow[r, equal] & W_a
\end{tikzcd}$$

Proof. We wish to construct a $\widetilde{W}$–equivariant section s to the restriction to $Q^\vee$ of the extension

$$1 \to Z/Z_c \to \widetilde{W_a}^{\mathrm{red}} \to W_a \to 1$$

Identify for this purpose $\widetilde{W_a}^{\mathrm{red}}$ with its image inside $N(H \times \mathbb{C}^*) \cap LG$ by using Proposition 8.9. We claim that there exists $x \in \mathbb{C}$ such that, for any $\lambda \in Q^\vee$, $\exp_H(x\lambda) \cdot z^\lambda$ lies in $\widetilde{W_a}^{\mathrm{red}}$. It is then clear that $s(\lambda) = \exp_H(x\lambda) \cdot z^\lambda$ yields the required section.

Let $\alpha_i^\vee$ be a short simple coroot and $w \in W$ an element such that $\theta^\vee = w\alpha_i^\vee$. Let $\widetilde{w} \in \widetilde{W}$ be a lift of w and lift $\theta^\vee \in Q^\vee$ to

$$\tau^{\theta^\vee} = \widetilde{s}_0 \, \widetilde{w} \widetilde{s}_i \widetilde{w}^{-1} \in \widetilde{W_a}^{\mathrm{red}}$$

Let $e_\theta \in \mathfrak{g}_\theta, f_\theta \in \mathfrak{g}_{-\theta}$ be root vectors such that $[e_\theta, f_\theta] = \theta^\vee$ and denote by $\rho_{\theta^\vee} : SL_2 \to G$ the embedding whose differential maps $e, f, h \in \mathfrak{sl}_2$ to $e_\theta, f_\theta, \theta^\vee$. We may assume e_θ, f_θ chosen so that $\widetilde{s}_0$ is of the form

$$\exp(f_\theta \otimes z)\exp(-e_\theta \otimes z^{-1})\exp(f_\theta \otimes z) = \rho_{\theta^\vee}(\begin{pmatrix} 0 & -z^{-1} \\ z & 0 \end{pmatrix})$$

Since $\widetilde{w}\widetilde{s}_i\widetilde{w}^{-1} \in N_\theta \setminus H_\theta$ is necessarily of the form

$$\exp(te_\theta)\exp(-t^{-1}f_\theta)\exp(te_\theta) = \rho_{\theta^\vee}(\begin{pmatrix} 0 & t \\ -t^{-1} & 0 \end{pmatrix})$$

for some $t \in \mathbb{C}^*$, we find that

$$\tau^{\theta^\vee} = \rho_{\theta^\vee}(\begin{pmatrix} (tz)^{-1} & 0 \\ 0 & tz \end{pmatrix}) = \exp(x\theta^\vee) \cdot z^{\theta^\vee}$$

with $x = -\ln(t)$ which proves our claim for $\lambda = \theta^\vee$. Let now $w \in W$ with lift $\widetilde{w} \in \widetilde{W}$, then

$$\widetilde{w}\tau_{\theta^\vee}\widetilde{w}^{-1} = \exp(xw(\theta^\vee)) \cdot z^{w\theta^\vee}$$

so that that $\exp(x\alpha_i^\vee) \cdot z^{\alpha_i^\vee} \in \widetilde{W_a}^{\text{red}}$ for any short coroot $\alpha_i^\vee$. Since the short coroots span $Q^\vee$, the claim holds for any $\lambda \in Q^\vee$.

Since $s(Q^\vee)$ is free abelian and $\widetilde{W}$ is finite, their intersection is trivial and the map $\widetilde{W} \ltimes Q^\vee \to \widetilde{W_a}^{\text{red}}$, $(\widetilde{w}, \lambda) \to \widetilde{w}s(\lambda)$ is injective. It is moreover surjective since Z/Z_c is generated by $\widetilde{s}_i^2$, $i = 1 \dots n$ and therefore lies in $\widetilde{W}$ ■

8.11.

Remark. Unlike $\widetilde{W_a}^{\text{red}}$, the (non–reduced) Tits extension $\widetilde{W_a}$ of W_a is not a semi–direct product in general. For example, for $\mathfrak{g} = \mathfrak{sl}_2$, with affine Cartan matrix

$$A = \begin{pmatrix} 2 & -2 \\ -2 & 2 \end{pmatrix}$$

$\widetilde{W_a}$ is generated by $\widetilde{s}_0$, $\widetilde{s}_1$ with relations $\widetilde{s}_i^4 = 1$,

$$\widetilde{s}_0\widetilde{s}_1^2\widetilde{s}_0^{-1} = \widetilde{s}_1^2(\widetilde{s}_0^2)^2 = \widetilde{s}_1^2 \quad \text{and} \quad \widetilde{s}_1\widetilde{s}_0^2\widetilde{s}_1^{-1} = \widetilde{s}_0^2$$

In particular, the group $Z \cong \mathbb{Z}_2^2$ generated by $\widetilde{s}_0^2, \widetilde{s}_1^2$ lies in the centre of $\widetilde{W_a}$. Any lift in $\widetilde{W_a}$ of the generator of $Q^\vee \cong \mathbb{Z}$ is of the form $\tau = z\widetilde{s}_0\widetilde{s}_1$, for some $z \in Z$ and gives rise a $\widetilde{W}$–equivariant section $Q^\vee \to \widetilde{W_a}$ if, and only if, $\widetilde{s}_1\tau\widetilde{s}_1^{-1} = \tau^{-1}$. Since $z = z^{-1}$ is central, such a section exists iff $\widetilde{s}_1(\widetilde{s}_0\widetilde{s}_1)\widetilde{s}_1^{-1} = \widetilde{s}_1^{-1}\widetilde{s}_0^{-1}$ and therefore iff $\widetilde{s}_1^2\widetilde{s}_0^2 = 1$ which holds in $\widetilde{W_a}^{\text{red}}$ but not in $\widetilde{W_a}$.

Remark. The section $Q^\vee \to \widetilde{W_a}^{\text{red}}$ constructed in Theorem 8.10 does not in general coincide with that obtained from the canonical section $Q^\vee \to B_a$ [Mc, §3.2–3.3]. For example, for $\mathfrak{g} = \mathfrak{sl}_3$, the canonical lift of $\theta^\vee \in Q^\vee$ in B_a is $T^{\theta^\vee} = S_0S_1S_2S_1$. When regarded as an element $\tau^{\theta^\vee}$ of $\widetilde{W_a}^{\text{red}}$ this does not give rise to a $\widetilde{W}$–equivariant section since $\operatorname{Ad}(\widetilde{s}_\theta)\tau^{\theta^\vee} \neq (\tau^{\theta^\vee})^{-1}$ where $\widetilde{s}_\theta = \widetilde{s}_1\widetilde{s}_2\widetilde{s}_1$ is a lift in $\widetilde{W}$ of the reflection s_θ. Indeed, $\operatorname{Ad}(\widetilde{s}_2)\widetilde{s}_1^2 = \widetilde{s}_1^2\widetilde{s}_2^2$ in $\widetilde{W}$, so that

$$\widetilde{s}_\theta^2 = \widetilde{s}_1 \operatorname{Ad}(\widetilde{s}_2)(\widetilde{s}_1^2)\widetilde{s}_2^2\widetilde{s}_1 = \widetilde{s}_1^3\widetilde{s}_2^4\widetilde{s}_1 = 1$$

Thus, since $\tau^{\theta^\vee} = \widetilde{s}_0\widetilde{s}_\theta$,

$$\operatorname{Ad}(\widetilde{s}_\theta)\tau^{\theta^\vee} = \widetilde{s}_\theta\widetilde{s}_0 \qquad \text{while} \qquad (\tau^{\theta^\vee})^{-1} = \widetilde{s}_\theta\widetilde{s}_0^{-1}$$

which are different elements of $\widetilde{W_a}^{\text{red}}$ by (8.6).

Acknowledgments

I am grateful to Raphaël Rouquier for pointing me towards the Yangian at the early stages of this project. I am also grateful to Sachin Gautam for spotting a gap in my initial proof of Theorem 2.5 and to the referee for a very careful reading of this manuscript. This paper was begun at the summer home of Pénélope Riboud and Romain Graziani. It is a pleasure to thank them for their warm and friendly hospitality, and their daughter Léonore for a number of inspiring comments.

References

[Br] A. Broer, *The sum of generalized exponents and Chevalley's restriction theorem for modules of covariants*, Indag. Math. (N.S.) **6** (1995), 385–396.

[BK] J. Brundan, A. Kleshchev, *Parabolic presentations of the Yangian $Y(\mathfrak{gl}_n)$*, Comm. Math. Phys. **254** (2005), 191–220.

[Ch1] I. Cherednik, *Affine extensions of Knizhnik–Zamolodchikov equations and Lusztig's isomorphisms*, Special functions (Okayama, 1990), 63–77, ICM-90 Satell. Conf. Proc., Springer, 1991.

[Ch2] I. Cherednik, *A unification of Knizhnik–Zamolodchikov and Dunkl operators via affine Hecke algebras*, Invent. Math. **106** (1991), no. 2, 411–431.

[Ch3] I. Cherednik, *Lectures on Knizhnik–Zamolodchikov equations and Hecke algebras*, Quantum many–body problems and representation theory, 1–96, MSJ Mem., **1**, Math. Soc. Japan, 1998.

[DCP] C. De Concini, C. Procesi, *On the geometry of toric arrangements*, Transform. Groups **10** (2005), 387–422.

[Dr1] V. G. Drinfeld, *Hopf algebras and the quantum Yang–Baxter equation*, Soviet Math. Dokl. **32** (1985), 254–258.

[Dr2] V. G. Drinfeld, *A new realization of Yangians and of quantum affine algebras*, Soviet Math. Dokl. **36** (1988), 212–216.

[Dr3] V. G. Drinfeld, *Degenerate affine Hecke algebras and Yangians*, Functional Anal. Appl. **20** (1986), 62–64.

[FMTV] G. Felder, Y. Markov, V. Tarasov, and A. Varchenko, *Differential Equations Compatible with KZ Equations*, Math. Phys. Anal. Geom. **3** (2000) 139–177.

[FR] I. B. Frenkel, N. Yu. Reshetikhin, *Quantum affine algebras and holonomic difference equations*, Comm. Math. Phys. **146** (1992), 1–60.

[GTL1] S. Gautam, V. Toledano Laredo, *Yangians and quantum affine algebras*, in preparation.

[GTL2] S. Gautam, V. Toledano Laredo, in preparation.

[Ka] V. G. Kac, *Infinite–Dimensional Lie Algebras*, 3rd Edition, Cambridge University Press, 1990.

[Ki] A. A. Kirillov Jr., *Lectures on affine Hecke algebras and Macdonald's conjectures*, Bull. Amer. Math. Soc. (N.S.) **34** (1997), 251–292.

[KR] A. N. Kirillov, N. Reshetikhin, *q–Weyl Group and a Multiplicative Formula for Universal R–Matrices*, Comm. Math. Phys. **134** (1990), 421–431.

[Ko] T. Kohno, *Integrable connections related to Manin and Schechtman's higher braid groups*, Illinois J. Math. **34** (1990), 476–484.

[Lo] E. Looijenga, *Arrangements, KZ systems and Lie algebra homology*, Singularity theory (Liverpool, 1996), 109–130, London Math. Soc. Lecture Note Ser., 263, Cambridge Univ. Press, 1999.

[Lu] G. Lusztig, *Affine Hecke algebras and their graded version*, J. Amer. Math. Soc. **2** (1989), 599–635.

[Mc] I. G. Macdonald, *Affine Hecke algebras and orthogonal polynomials*. Cambridge Tracts in Mathematics, 157. Cambridge University Press, Cambridge, 2003.

[Mo] A. Molev, *Yangians and classical Lie algebras*. Mathematical Surveys and Monographs, 143. American Mathematical Society, 2007.

[MTL] J. J. Millson, V. Toledano Laredo, *Casimir operators and monodromy representations of generalised braid groups*, Transform. Groups **10** (2005) 217–254.

[NO] M. Nazarov, G. Olshanski, *Bethe subalgebras in twisted Yangians*, Comm. Math. Phys. **178** (1996), 483–506.

[NVD] Nguyêñ Viêt Dũng, *The fundamental groups of the spaces of regular orbits of the affine Weyl groups*, Topology **22** (1983), 425–435.

[PS] A. Pressley, G. Segal, *Loop groups.* Oxford Mathematical Monographs. Oxford University Press, 1986.

[Re] M. Reeder, *Exterior powers of the adjoint representation*, Canad. J. Math. **49** (1997), 133–159.

[Re2] M. Reeder, *Zero weight spaces and the Springer correspondence*, Indag. Math. (N.S.) **9** (1998), 431–441.

[RS] A. Ram, A. Shepler, *Classification of graded Hecke algebras for complex reflection groups*, Comment. Math. Helv. **78** (2003), 308–334.

[So] Y. S. Soibelman, *Algebra of Functions on a Compact Quantum Group and its Representations*, Leningrad Math. J. **2** (1991), 161–178.

[Va] M. Varagnolo, *Quiver varieties and Yangians*, Lett. Math. Phys. **53** (2000), 273–283.

[TV] V. Tarasov, A. Varchenko, *Dynamical differential equations compatible with rational qKZ equations*, Lett. Math. Phys. **71** (2005), 101–108.

[Ti] J. Tits, *Normalisateurs de tores. I. Groupes de Coxeter étendus*, J. Algebra **4** (1966), 96–116.

[TL1] V. Toledano Laredo, *A Kohno–Drinfeld theorem for quantum Weyl groups*, Duke Math. J. **112** (2002), 421–451.

[TL2] V. Toledano Laredo, *Flat Connections and Quantum Groups*, Acta Appl. Math. **73** (2002), 155–173.

[TL3] V. Toledano Laredo, *Quasi–Coxeter algebras, Dynkin diagram cohomology and quantum groups*, Int. Math. Res. Pap. **2008**, Art. ID rpn009, 167 pp.

[vdL] H. van der Lek, *Extended Artin groups*. Singularities, Part 2 (Arcata, Calif., 1981), 117–121, Proc. Sympos. Pure Math., 40, Amer. Math. Soc., 1983.

Department of Mathematics, Northeastern University, 567 Lake Hall, 360 Huntington Avenue, Boston, MA 02115.

E-mail address: `V.ToledanoLaredo@neu.edu`